

Alpha-Vacua, Black Holes, and AdS/CFT

Andrew Chamblin¹ and Jeremy Michelson^{2,*}

¹*Somewhere in Bush Country, specifically,*

*Department of Physics
102 Natural Sciences Bldg.
University of Louisville
Louisville, KY 40292
USA*

²*Department of Physics and Astronomy*

*University of Kentucky
600 Rose Street
Lexington, KY 40506
USA*

Abstract

The Schwarzschild, Schwarzschild-AdS, and Schwarzschild-de Sitter solutions all admit freely acting discrete involutions which commute with the continuous symmetries of the spacetimes. Intuitively, these involutions correspond to the antipodal map of the corresponding spacetimes. In analogy with the ordinary de Sitter example, this allows us to construct new vacua by performing a Mottola-Allen transform on the modes associated with the Hartle-Hawking, or Euclidean, vacuum. These vacua are the “alpha”-vacua for these black holes. The causal structure of a typical black hole may ameliorate certain difficulties which are encountered in the case of de Sitter α -vacua. For Schwarzschild-AdS black holes, a Bogoliubov transformation which mixes operators of the two boundary CFT’s provides a construction of the dual CFT α -states. Finally, we analyze the thermal properties of these vacua.

*Electronic address: jeremy@pa.uky.edu; Current address: Department of Physics; The Ohio State University; 1040 Physics Research Building; 191 West Woodruff Avenue; Columbus, Ohio 43210-1117; USA

Contents	
I. Introduction	3
Note:	4
II. The Discrete Involutions of Schwarzschild	4
III. Geometry of a General, Neutral, Spherically Symmetric Black Hole	7
IV. α-Vacua	8
A. Modes	8
B. Green Functions	10
C. Example: The BTZ Black Hole	13
D. The Hartle-Hawking Vacuum	15
1. Propagators	15
2. Thermality	17
E. Thermality of α -Vacua	18
V. (No) Objections to α-Vacua	18
A. Causality	19
B. Poles	20
C. Thermality	20
D. Pinch Singularities	22
E. Uniqueness of Black Hole Vacua	24
VI. CFT α-States	25
VII. AdS/CFT	27
A. Populations	27
B. Propagators	28
C. Entropy	29
Acknowledgments	30
A. Derivation of the CFT Density Matrix	30
B. Strings on Rindler Space	31
1. The Setup and Vertex Operators	31
2. Rindler Space α -Vacua	33
3. No Pinch Singularities in One-loop Diagrams	34
a. Minkowski space diagrams	34
b. The Three-Point Function	35
References	37

I. INTRODUCTION

The α -vacua [1, 2] of a scalar field in the de Sitter (dS) spacetime have generated considerable interest (for example, Refs. [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]). At first glance, these vacua appear to be highly unphysical because they exhibit rather sick behaviour in the UV. However, the idea is that they still may be important for cosmology because they may correspond to certain quantum gravity effects which have yet to be fully understood. For example, it has been suggested that the value of α may be imprinted in the cosmic microwave background. Discussions include [4, 14, 15, 16, 17, 18, 19] and other relevant papers are [20, 21, 22, 23, 24]. α -Vacua for spin 1/2 fields were discussed in [25].

Although α -vacua have an unorthodox short distance behaviour, they are natural in that they preserve all of the de Sitter isometries. In this sense, there is no reason to prefer one value of α over any other value. That said, the literature [4, 5, 6, 7, 8, 9] contains many reasons, besides short distance physics, for preferring the Euclidean vacuum obtained by analytic continuation from Euclidean dS, to the other α -vacua.

It has been suggested that some of the objectionable characteristics of α -vacua can be avoided by changing the time ordering prescription of operators, in an α -dependent way.[10, 11] However,[8] the path integral naturally picks out the usual time ordering prescription for the Feynman propagator. It is therefore unclear to us what a change in the time ordering would mean, or how it would be natural. Thus we stick with the usual meaning of time ordering.

In this work, we introduce the notion of α -vacua for black hole spacetimes. We start by discussing Schwarzschild, but in fact the construction works in fair generality. The black hole horizon allows us to evade many of the objections that have been raised in the context of de Sitter α -vacua. (We restrict attention to neutral, nonrotating black holes and do not address whether α -vacua can be defined for black holes whose singularities are not spacelike.) In particular, we show that Schwarzschild-AdS solutions admit α -vacua.

These vacua could have profound implications for Schwarzschild-AdS and the AdS/CFT correspondence. This is because there are two distinct boundary components, and therefore two distinct theories, which encode the bulk geometry of Schwarzschild-AdS. The α -vacua explicitly “correlate” points in the two different theories. Maldacena [26] has suggested that it might be possible to realize the entropy of Schwarzschild-AdS as an entanglement entropy between the two theories.

We describe how the α -vacua can be constructed not only on the AdS side but also on the CFT side. On the CFT side, one essentially has to perform a Bogoliubov transformation that mixes operators of the two CFTs, while continuing to trace over a single *boundary*. (This prescription necessarily differs from the prescription [3] used in dS/CFT.) We show that this modifies the CFT in accord with the AdS—and also introduces α -dependence into the entropy, at least for the naïve definition of the entropy.

In [27, 28] α -vacua were constructed for certain asymptotically AdS black holes—not those considered in this paper—from the de Sitter α -vacua by using the de Sitter slicing of AdS. In particular, [28] showed that most α -vacua defined in this way are rendered ill-defined by the black hole singularity; of this family of vacua, only the Euclidean vacuum is well-defined. Our construction of α -vacua is different, and does not rely on a foliation of the spacetime by de Sitter slices.

This paper is organized as follows. In §II we discuss the geometry of Schwarzschild, and find a freely acting antipodal symmetry which allows us to define α -vacua. We extend this

to a much larger class of spacetimes, including Schwarzschild-AdS, in §III. α -vacua are then defined and examined in §IV. Objections to α -vacua for de Sitter are reviewed in §V and then rebutted for the spacetimes considered in this paper. Comments therein regarding string theory amplitudes for α -vacua in Rindler space are elaborated on in Appendix B. The CFT-dual states to α -vacua are described in §VI. Consequences of AdS/CFT are then described in §VII. We ensure that the CFT states, that we propose are dual to AdS α -vacua, do indeed reproduce AdS propagators and particle production. We close with the observation that black hole entropy is α -dependent.

Note:

This project was initiated about two years ago with §II, which resulted from a conversation between Andrew, S. Das and A. Shapere while Andrew was driving them in his Land Rover, and Andrew was actively involved with all of the essential part of the work. Unfortunately, Andrew passed away while this work was waiting to be finalized. J.M. thanks Andrew's friends, family and colleagues for the opportunity to have presented this work at the Andrew Chamblin Memorial Symposium in March 2006 at the University of Louisville. J.M. also apologizes for the delay in producing this last paper of Andrew's—and thanks those who encouraged J.M. to get Andrew's paper into final form. Andrew will be missed.

II. THE DISCRETE INVOLUTIONS OF SCHWARZSCHILD

To begin, let $(\mathcal{M}, g)$ denote the Schwarzschild spacetime. We are interested in constructing discrete involutive isometries which will act on $\mathcal{M}$. In particular, we are interested in the actions of time and space inversion. Since we wish to consider the actions of these inversions on the maximally extended spacetime, it is most natural to use Kruskal coordinates [29], which cover the entire manifold. We therefore begin with a review of the relation of these coordinates to the usual Schwarzschild coordinates (which cover only part of the maximal extension).

Let (t, r, θ, ϕ) denote the Schwarzschild coordinates so that the metric reads

$$ds^2 = -\left(1 - \frac{2m}{r}\right) dt^2 + \frac{dr^2}{\left(1 - \frac{2m}{r}\right)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (2.1)$$

Next introduce null coordinates u and v such that

$$du = dt - \frac{dr}{\left(1 - \frac{2m}{r}\right)}, \quad dv = dt + \frac{dr}{\left(1 - \frac{2m}{r}\right)}, \quad (2.2)$$

or, integrating

$$u = t - r - 2m \log(r - 2m), \quad v = t + r + 2m \log(r - 2m). \quad (2.3)$$

Now form the coordinates U and V by exponentiating

$$U = -e^{-\frac{u}{4m}}, \quad V = e^{\frac{v}{4m}}, \quad (2.4)$$

Then one finds that the coordinates T and Z defined by

$$T = \sinh\left(\frac{t}{4m}\right) e^{\frac{r}{4m}} \sqrt{r-2m}, \quad Z = \cosh\left(\frac{t}{4m}\right) e^{\frac{r}{4m}} \sqrt{r-2m}, \quad (2.5)$$

satisfy the simple algebraic relations

$$U = T - Z, \quad V = T + Z, \quad (2.6)$$

i.e., U and V are advanced and retarded null coordinates relative to T and Z . One checks that in these coordinates the metric assumes the form

$$ds^2 = e^{-\frac{r}{2m}} \left(16 \frac{m^2}{r}\right) (-dT^2 + dZ^2) + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (2.7)$$

Using the coordinates (T, Z, θ, ϕ) , we define total time inversion by the map

$$R_T : (T, Z, \theta, \phi) \longrightarrow (-T, Z, \theta, \phi), \quad (2.8)$$

and likewise space inversion is given by

$$R_Z : (T, Z, \theta, \phi) \longrightarrow (T, -Z, \theta, \phi). \quad (2.9)$$

Of course, neither of these involutions acts freely (they both have fixed points). To obtain a free action, we need to take a product with some other map which *is* freely acting. Such a map, which we denote P , is given as

$$P : (T, Z, \theta, \phi) \longrightarrow (T, Z, \pi - \theta, \phi + \pi). \quad (2.10)$$

Thus, we can construct the following four freely acting involutions on $\mathcal{M}$: P, PR_T, PR_Z and $PR_Z R_T$. We claim that all of these involutions extend to the corresponding Euclidean instanton (the “cigar”). Before addressing the Riemannian issue, however, we need to first determine which of these maps commutes with the continuous symmetries of the spacetime. This was actually studied by Gibbons [30], who found that only the map $J = PR_Z R_T$ commutes with the continuous symmetries of $\mathcal{M}$. We shall refer to this antipodal map as the “CPT” operator. If we consider the *quotient* manifold

$$\mathcal{M}_J = \mathcal{M}/J \quad (2.11)$$

then it is straightforward to show that $\mathcal{M}_J$ is asymptotically flat, is *not* time orientable, and is space orientable. In order to see how these involutions extend to the Euclidean section, it is useful to write complexified Schwarzschild as an algebraic variety in $\mathbb{C}^7$ in the usual fasion. Explicitly, let $\{Z^i \mid i = 1, \dots, 7\}$ be coordinates on $\mathbb{C}^7$, so that in terms of Schwarzschild coordinates (which cover only a subset of the variety), we have

$$\begin{aligned} Z^1 &= r \sin \theta \cos \phi, & Z^4 &= -2m\sqrt{\frac{2m}{r}} + 4m\sqrt{\frac{r}{2m}}, & Z^6 &= 4m\sqrt{1 - \frac{2m}{r}} \cosh\left(\frac{t}{4m}\right), \\ Z^2 &= r \sin \theta \sin \phi, & & & & \\ Z^3 &= r \cos \theta, & Z^5 &= 2\sqrt{3}m\sqrt{\frac{2m}{r}}, & Z^7 &= 4m\sqrt{1 - \frac{2m}{r}} \sinh\left(\frac{t}{4m}\right). \end{aligned} \quad (2.12)$$

With the coordinates as in (2.7), it turns out that complexified Schwarzschild ($\mathcal{M}_{\mathbb{C}}$) is given as the algebraic variety determined by the three polynomials

$$\begin{aligned} (Z^6)^2 - (Z^7)^2 + \frac{4}{3}(Z^5)^2 &= 16m^2, \\ [(Z^1)^2 + (Z^2)^2 + (Z^3)^2] (Z^5)^4 &= 576m^6, \\ \sqrt{3}Z^4Z^5 + (Z^5)^2 &= 24m^2. \end{aligned} \quad (2.13)$$

The Lorentzian section ($\mathcal{M} = \mathcal{M}^L$) and the Riemannian section ($\mathcal{M}^R$) are determined by finding certain anti-holomorphic involutions acting on the above variety which stabilise either $\mathcal{M}^L$ or $\mathcal{M}^R$; that is, we find maps

$$J_L : \mathcal{M}_{\mathbb{C}} \longrightarrow \mathcal{M}_{\mathbb{C}}, \quad J_R : \mathcal{M}_{\mathbb{C}} \longrightarrow \mathcal{M}_{\mathbb{C}}, \quad (2.14)$$

such that J_L leaves $\mathcal{M}^L \subset \mathcal{M}_{\mathbb{C}}$ invariant:

$$J_L(\mathcal{M}^L) = \mathcal{M}^L, \quad (2.15)$$

and such that J_R leaves $\mathcal{M}^R \subset \mathcal{M}_{\mathbb{C}}$ invariant:

$$J_R(\mathcal{M}^R) = \mathcal{M}^R. \quad (2.16)$$

As described in [31], J_L restricted to $\mathcal{M}^L$ is an anti-holomorphic version of time reversal. J_R is the map given by reflection through the $\tau = 0$ (where $\tau = it$) three-surface in the “cigar” instanton (*i.e.*, $\tau = 0$ is the “Einstein Rosen bridge” three-surface Σ , with topology $S^2 \times \mathbb{R}$). Since the surfaces $t = 0$ and $\tau = it = 0$ correspond to the surface $Z^7 = 0$, we see that $\mathcal{M}^L$ and $\mathcal{M}^R$ intersect precisely along this Einstein Rosen bridge. Explicitly, we can realise the two maps J_L and J_R as follows:

$$\begin{aligned} J_L : (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6, Z^7) &\longrightarrow (\bar{Z}^1, \bar{Z}^2, \bar{Z}^3, \bar{Z}^4, \bar{Z}^5, \bar{Z}^6, \bar{Z}^7), \\ J_R : (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6, Z^7) &\longrightarrow (\bar{Z}^1, \bar{Z}^2, \bar{Z}^3, \bar{Z}^4, \bar{Z}^5, \bar{Z}^6, -\bar{Z}^7). \end{aligned} \quad (2.17)$$

Comparing these explicit formulae for J_L and J_R with the coordinates in (2.7), we see that J_R is thus obtained from J_L by the transformation $t \longrightarrow \tau = it$.

What we want to do now is show how the maps R_T, R_Z and P acting on $\mathcal{M}^L$, and likewise their Euclidean counterparts acting on $\mathcal{M}^R$, are actually just the *restrictions* to $\mathcal{M}^L$ and $\mathcal{M}^R$ of certain *holomorphic* involutions acting on $\mathcal{M}_{\mathbb{C}}$. Of course, once we notice that our complex coordinates Z^6 and Z^7 are (up to a scaling) actually our Kruskal coordinates Z and T , it is easy to see that the “big” involutions, $\mathcal{R}_Z$ and $\mathcal{R}_T$ (which restrict to R_Z and R_T on $\mathcal{M}^L$) are given by

$$\mathcal{R}_Z : (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6, Z^7) \longrightarrow (Z^1, Z^2, Z^3, Z^4, Z^5, -Z^6, Z^7), \quad (2.18)$$

$$\mathcal{R}_T : (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6, Z^7) \longrightarrow (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6, -Z^7). \quad (2.19)$$

Clearly, these maps are holomorphic, and since they commute with both J_L and J_R , they restrict to well-defined involutions on $\mathcal{M}^L$ and $\mathcal{M}^R$. Thus, $\mathcal{R}_Z|_{\mathcal{M}^L} = R_Z$ and $\mathcal{R}_T|_{\mathcal{M}^L} = R_T$. For the maps restricted to the Riemannian section, we shall write

$$\mathcal{R}_Z|_{\mathcal{M}^R} = \bar{R}_Z : \mathcal{M}^R \longrightarrow \mathcal{M}^R \quad (2.20)$$

$$\mathcal{R}_T|_{\mathcal{M}^R} = \bar{R}_T : \mathcal{M}^R \longrightarrow \mathcal{M}^R \quad (2.21)$$

In terms of local coordinates on $\mathcal{M}^R$, these reflections take the form

$$\bar{R}_Z : \tau \longrightarrow -\tau + 4\pi m, \quad \bar{R}_T : \tau \longrightarrow -\tau \quad (2.22)$$

(r, θ , and ϕ are left invariant by both these maps). Thus, we see that $\bar{R}_T$ reflects in imaginary time whereas $\bar{R}_Z$ corresponds to rotating through half a period in imaginary time.

Finally, we obtain the involution $\bar{P}$ on $\mathcal{M}^R$ by restricting to $\mathcal{M}^R$ the following map on $\mathcal{M}^{\mathbb{C}}$:

$$\mathcal{P} : (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6, Z^7) \longrightarrow (-Z^1, -Z^2, -Z^3, Z^4, Z^5, Z^6, Z^7). \quad (2.23)$$

Now that we have made sense of how to extend our discrete isometries R_Z, R_T and P from $\mathcal{M}$ to $\mathcal{M}^R$, we can act on the Hartle-Hawking modes of the Kruskal manifold with these involutions. This allows us to apply the Mottola-Allen transformation and so obtain α -vacua. It should be obvious that one may obtain these Euclidean involutions for any spherically symmetric black hole spacetime, and so we now turn our attention to the more general case.

III. GEOMETRY OF A GENERAL, NEUTRAL, SPHERICALLY SYMMETRIC BLACK HOLE

Following [32], we consider the $(d+1)$ -dimensional spacetime with metric

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_{d-1}^2, \quad (3.1)$$

where $f(r)$ is a monotonically increasing (for positive r) function with a singularity at $r = 0$ and a simple zero at $r = r_+ > 0$. The assumed monotonicity implies that $r = r_+$ is the only real, positive zero. The singularity at $r = 0$ is assumed to give a spacetime singularity, and so the spacetime cannot be extended beyond $r = 0$. Examples are Schwarzschild [$f(r) = 1 - \frac{\omega_d m}{r^{d-2}}$] and Schwarzschild-AdS [$f(r) = \frac{r^2}{\ell^2} + 1 - \frac{\omega_d m}{r^{d-2}}$] where ℓ is the length-scale associated with the cosmological constant, and m is the properly normalized mass of the black hole, when $\omega_d = \frac{16\pi G_N}{(d-1)\text{Vol}(S^{d-1})}$, with G_N the Newton constant.

There is a coordinate singularity (horizon) at $f(r) = 0$, *i.e.* $r = r_+$. We extend the coordinates past the horizon, and eventually define Kruskal coordinates [29] by first defining the tortoise coordinates r^* by

$$r^*(r) = \int_0^r \frac{dr'}{f(r')} + \frac{\pi i}{f'(r_+)}. \quad (3.2)$$

The rationale for the additional, imaginary constant is as follows.[32] We want the coordinate change to be well-defined in the region of overlap with the original coordinates, namely $r > r_+$. But the integration from $r = 0$ passes through the pole at $r = r_+$ and contributes an imaginary part to the integral. This has been explicitly cancelled (with the contour going below the pole).

Next define the double-null coordinates

$$u = t - r^*, \quad v = t + r^*. \quad (3.3)$$

The metric reads

$$ds^2 = -f(r)dudv + r^2 d\Omega_{d-1}^2, \quad (3.4)$$

where, of course, r is now considered a function of u and v . Since $f(r)$ was assumed to be monotonic, $r > 0$ is, in fact, a single valued function of $v - u$. Clearly, however, this metric is still singular at $r = r_+$.

With the conformal transformation

$$U = -e^{-\frac{f'(r_+)}{2}u} = T - Z, \quad V = e^{\frac{f'(r_+)}{2}v} = T + Z, \quad (3.5)$$

the metric becomes

$$\begin{aligned} ds^2 &= -\frac{4}{f'(r_+)^2} f(r) e^{-f'(r_+)r^*} dU dV + r^2 d\Omega_{d-1}^2 \\ &= \frac{4}{f'(r_+)^2} f(r) e^{-f'(r_+)r^*} (-dT^2 + dZ^2) + r^2 d\Omega_{d-1}^2, \end{aligned} \quad (3.6)$$

and we have the usual identities

$$UV = T^2 - Z^2 = -e^{f'(r_+)r^*}, \quad \tanh^{-1} \frac{T}{Z} = \frac{f'(r_+)}{2} t. \quad (3.7)$$

The metric is now smooth at $r = r_+$, and therefore the U, V coordinates extend to the singularity at $r = 0$. These are the Kruskal coordinates from which one can easily obtain the Penrose diagram. The shape of the Penrose diagram depends on $r^*(r = \infty)$, as shown in Figure 1.¹

The exterior metric (3.1) clearly has spherical and time-translation symmetry. These clearly extend to the entire spacetime, although the isometry corresponding to time translation is null on the horizons, and spacelike inside the black and white holes. Explicitly,

$$\frac{\partial}{\partial t} = \frac{f'(r_+)}{2} \left(-U \frac{\partial}{\partial U} + V \frac{\partial}{\partial V} \right) = \frac{f'(r_+)}{2} \left(T \frac{\partial}{\partial Z} + Z \frac{\partial}{\partial T} \right). \quad (3.8)$$

IV. α -VACUA

A. Modes

Recall the definitions (2.8), (2.9), (2.10). Although R_T and R_Z are symmetries of the spacetime—note that r^* and therefore r is invariant under each map—neither preserves the time-like Killing vector $\frac{\partial}{\partial t}$. The combination $R_Z R_T$ preserves all the symmetries of the spacetime—at least for generic $f(r)$ —but has a fixed point at $T = Z = 0$. The product $PR_Z R_T$ acts freely and preserves all the (generic) symmetries of the spacetime. We refer to this as the antipodal map, $\mathcal{P}_A$,

$$\mathcal{P}_A = PR_Z R_T. \quad (4.1)$$

¹ The authors of [32] use rescaled Penrose coordinates relative to us, and so obtain bowed singularities instead of bowed asymptotics. However, this rescaling is not possible for asymptotically flat spacetimes, so we have consistently drawn bowed asymptotics.




FIG. 1: Penrose diagrams of the various types of spacetimes. The exterior and interior regions have been labelled in the usual way. The direction of time, Eq. (3.8), in each region is also indicated. (a) $r^*(r = \infty) = \infty$ (e.g. asymptotically flat spacetimes.) (b) $r^*(r = \infty) = 0$ (e.g. the BTZ black hole). (c) $0 < r^*(r = \infty) < \infty$. Large mass AdS_5 -Schwarzschild is the example drawn, for which $r^*(r = \infty) = \pi$. (d) $r^*(r = \infty) < 0$. The figure is for $r^*(r = \infty) = -\pi$.

Observe that the antipodal map connects points in opposite exterior regimes—that is, points for which time has opposite orientation. The antipodal point of $x = (T, Z, \Omega)$ is denoted $x_A = (-T, -Z, P\Omega)$.

The wave equation $(-\square + \mu^2)\phi = 0$ for a “free” scalar field ϕ of mass μ reads²

$$\frac{UVf'(r_+)^2}{8f(r)r^{d-1}} \left[\frac{\partial}{\partial U} r^{d-1} \frac{\partial}{\partial V} + \frac{\partial}{\partial V} r^{d-1} \frac{\partial}{\partial U} \right] \phi - \frac{1}{r^2} \nabla_{S^{d-1}}^2 \phi + \mu^2 \phi = 0, \quad r = r(UV). \quad (4.2)$$

This is obviously solved by expanding in spherical harmonics $Y_{n,s^{(n)}}(\Omega)$ on the S^{d-1} for which

$$\nabla_{S^{d-1}}^2 Y_{n,s^{(n)}}(\Omega) = -n(n + d - 2) Y_{n,s^{(n)}}(\Omega). \quad (4.3)$$

Here $s^{(n)}$ denotes the remaining quantum numbers of the spherical harmonics.

² For simplicity, we assume either no coupling of the field to the scalar curvature, or else that such a coupling can be made, and is, part of the definition of μ to the linearized level. The latter assumption is equivalent to demanding that the spacetime have constant scalar curvature, which, of course, restricts f , but is consistent with demanding a vacuum solution of Einstein’s equations with or without cosmological constant.

We choose a standard [2, 3] nonstandard (not a typo) basis of spherical harmonics for which

$$Y_{n,s(n)}(P\Omega) = Y_{n,s(n)}(\Omega)^*, \quad (4.4)$$

with the asterisk denoting complex conjugation. Since the equation of motion (4.2) is clearly invariant under complex conjugation and under the antipodal map, we use these spherical harmonics to choose our modes such that

$$\phi_{\kappa,n,s(n)}(x_A) = \phi_{\kappa,n,s(n)}(x)^*, \quad (4.5)$$

where κ is the remaining quantum number(s) needed to specify the solution, and is (presumably) related to the frequency of the mode. The Klein-Gordon inner product implies that if $\phi_{\kappa,n,s(n)}(x)$ is a “positive frequency” mode then $\phi(x)_{\kappa,n,s(n)}^*$ is “negative frequency”. This is, however, consistent with the choice (4.5) as the antipodal map includes time reversal.

The mode expansion reads

$$\phi(x) = \sum_{\kappa,n,s(n)} \left[a_{\kappa,n,s(n)} \phi_{\kappa,n,s(n)}(x) + a_{\kappa,n,s(n)}^\dagger \phi_{\kappa,n,s(n)}^*(x) \right], \quad (4.6)$$

Properly normalized modes under the Klein-Gordon inner-product imply canonical commutation relations for the operators,

$$\left[a_{\kappa,n,s(n)}, a_{\kappa',n',s'(n')}^\dagger \right] = \delta_{\kappa,\kappa'} \delta_{n,n'} \delta_{s(n),s'(n')}, \quad (4.7)$$

where, for κ the sum and Kronecker- δ is understood as a an integration, possibly with discrete sum, and Dirac- δ , possibly with an additional Kronecker- δ , if κ takes continuum values, possibly with an additional discrete quantum number. This is understood without comment in the following.

B. Green Functions

The vacuum is defined to be annihilated by annihilation operators $a_{\kappa,n,s(n)}$. Green functions are then found by evaluating two-point functions. The Wightman function is

$$G_0^{(+)}(x, x') = \langle 0 | \phi(x) \phi(x') | 0 \rangle = \sum_{\kappa,n,s(n)} \phi_{\kappa,n,s(n)}(x) \phi_{\kappa,n,s(n)}^*(x'); \quad (4.8)$$

the Hadamard function is

$$G_0^{(1)}(x, x') = \langle 0 | \{ \phi(x), \phi(x') \} | 0 \rangle = G^{(+)}(x, x') + G^{(+)}(x', x); \quad (4.9)$$

the commutator function is

$$iD_0(x, x') = \langle 0 | [\phi(x), \phi(x')] | 0 \rangle = G^{(+)}(x, x') - G^{(+)}(x', x) = G^{(+)}(x, x') - G^{(+)}(x, x')^*; \quad (4.10)$$

and the Feynman propagator,

$$\begin{aligned} iG_0^F(x, x') &= \langle 0 | \mathcal{T}(\phi(x) \phi(x')) | 0 \rangle = \Theta(x, x') G_0^{(+)}(x, x') + \Theta(x', x) G_0^{(+)}(x', x) \\ &= \frac{1}{2} G_0^{(1)}(x, x') + \frac{1}{2} \epsilon(x, x') iD_0(x, x'), \end{aligned} \quad (4.11)$$

where $\mathcal{T}$ denotes time ordering and

$$\Theta(x, x') = \begin{cases} 1, & x \text{ to the future of } x', \\ \frac{1}{2}, & x, x' \text{ spacelike separated,} \\ 0, & \text{otherwise,} \end{cases} \quad \epsilon(x, x') = \begin{cases} 1, & x \text{ to the future of } x', \\ 0, & x, x' \text{ spacelike separated,} \\ -1, & x \text{ to the past of } x'. \end{cases} \quad (4.12)$$

In general, however, there is no natural choice of “positive frequency” modes and therefore no natural choice of vacuum; given a particular decomposition, one can always perform a Bogoliubov transformation to a new, nonequivalent, decomposition. However, there are natural choices in the following senses. First, if one demands that the vacuum respect the symmetries of the spacetime, then this restricts the choices of vacua. Indeed, it is well known that the Minkowski vacuum is unique. For de Sitter, there is a one (complex)-parameter family of symmetric vacua (α -vacua), obtained from the Bunch-Davies vacuum by a $\vec{k}$ -independent Bogoliubov transformation, where $\vec{k}$ is the spatial momentum in planar coordinates.[1, 2] Alternatively, one usually obtains a unique vacuum by analytic continuation from the Euclidean spacetime, assuming the latter exists; for de Sitter, this Hartle-Hawking vacuum coincides with the Bunch-Davies vacuum.

In Ref. [33], it is argued that it is the Boulware vacuum, and not the Hartle-Hawking vacuum, that is relevant to AdS/CFT for black hole backgrounds. However, it should be clear that most of our discussion does not depend on the details of the vacuum with which we start. One can also construct α -vacua starting with the Boulware vacuum, and our discussion will follow more-or-less identically.

Note that $T \rightarrow iT_E$ in (3.6) defines a perfectly well-defined Euclidean spacetime and $r \geq r_+$. $r = r_+$ is the origin $T_E = Z = 0$. Alternatively, $t \rightarrow it_E$ also gives a well-defined Euclidean spacetime with $r \geq r_+$. In particular, observe that U and V are periodic as $t \rightarrow t + \frac{4\pi i}{f'(r_+)}$; this suggests that an observer in the original (t, r) -coordinates sees a temperature $\frac{1}{\beta} = \frac{f'(r_+)}{4\pi}$.

It seems difficult to analyze the symmetries preserved by a general Bogoliubov transformation without having explicit formulas for modes. However, if we can write down Green functions solely in terms of geodesic distances, then we have certainly found an invariant vacuum. Our strategy is therefore to study the Green function following Hartle and Hawking [34]. This will not give us precise detailed information, but will give us enough analytic structure to study generic properties.

Once we have the Hartle-Hawking Green function, then we can define α -vacua following [2]. In terms of modes, we define

$$\tilde{\phi}_{\kappa, n, s(n)}(x) = \cosh \alpha \phi_{\kappa, n, s(n)}(x) + e^{i\gamma} \sinh \alpha \phi_{\kappa, n, s(n)}^*(x), \quad (4.13)$$

where α and γ are real.³ This defines new annihilation and creation operators

$$\phi(x) = \sum_{\kappa, n, s(n)} \left[\tilde{a}_{\kappa, n, s(n)} \tilde{\phi}_{\kappa, n, s(n)} + \tilde{a}_{\kappa, n, s(n)}^\dagger \tilde{\phi}_{\kappa, n, s(n)}^* \right], \quad (4.14)$$

³ Usually one uses β rather than γ , but that would confuse the phase with the inverse temperature. An alternate convention is $\tilde{\phi}_{\kappa, n, s(n)}(x) = N_\alpha \left(\phi_{\kappa, n, s(n)}(x) + e^\alpha \phi_{\kappa, n, s(n)}^*(x) \right)$, where $\text{Re } \alpha < 0$ and $N_\alpha = 1/\sqrt{1 - e^{\alpha + \alpha^*}}$. In other words, $e^{\alpha_{\text{alternate}}} = e^{i\gamma} \tanh \alpha_{\text{here}}$.

$$\begin{aligned}\tilde{a}_{\kappa,n,s(n)} &= \cosh \alpha a_{\kappa,n,s(n)} - e^{-i\gamma} \sinh \alpha a_{\kappa,n,s(n)}^\dagger, \\ \tilde{a}_{\kappa,n,s(n)}^\dagger &= \cosh \alpha a_{\kappa,n,s(n)}^\dagger - e^{i\gamma} \sinh \alpha a_{\kappa,n,s(n)}.\end{aligned}\tag{4.15}$$

The vacuum annihilated by $\tilde{a}_{\kappa,n,s(n)}$ is the α -vacuum $|\alpha\gamma\rangle$. It preserves the symmetries of the spacetime if $|0\rangle$ does. Specifically, because of the choice (4.5), we have, for the Wightman function for example,

$$\begin{aligned}G_{\alpha\gamma}^{(+)}(x, x') &= \langle \alpha\gamma | \phi(x) \phi(x') | \alpha\gamma \rangle = \cosh^2 \alpha G_0^{(+)}(x, x') + e^{i\gamma} \sinh \alpha \cosh \alpha G_0^{(+)}(x_A, x') \\ &\quad + e^{-i\gamma} \sinh \alpha \cosh \alpha G_0^{(+)}(x, x'_A) + \sinh^2 \alpha G_0^{(+)}(x_A, x'_A).\end{aligned}\tag{4.16}$$

Since the antipodal map commutes with the symmetries of the spacetime, and since $|0\rangle$ was assumed to preserve the symmetries, thereby implying that $G_0^{(+)}(x, x')$ respects the symmetries, it follows that $G_{\alpha\gamma}^{(+)}(x, x')$ respects the symmetries of the spacetime. Also, note that

$$G_0^{(+)}(x_A, x'_A) = G_0^{(+)}(x', x) = G_0^{(+)}(x, x')^*.\tag{4.17}$$

Therefore,

$$\begin{aligned}G_{\alpha\gamma}^{(1)}(x, x') &= G_{\alpha\gamma}^{(+)}(x, x') + G_{\alpha\gamma}^{(+)}(x', x), \\ &= (\cosh^2 \alpha + \sinh^2 \alpha) G_0^{(1)}(x, x') + e^{i\gamma} \sinh 2\alpha G_0^{(+)}(x_A, x') + e^{-i\gamma} \sinh 2\alpha G_0^{(+)}(x, x'_A), \\ &= \cosh 2\alpha G_0^{(1)}(x, x') + \cos \gamma \sinh 2\alpha G_0^{(1)}(x_A, x') - \sin \gamma \sinh 2\alpha D_0(x_A, x').\end{aligned}\tag{4.18}$$

Similarly,

$$iD_{\alpha\gamma}(x, x') = G_{\alpha\gamma}^{(+)}(x, x') - G_{\alpha\gamma}^{(+)}(x', x) = iD_0(x, x');\tag{4.19}$$

this also follows since the commutator of the field is a c-number, independent of the vacuum. Finally,

$$\begin{aligned}iG_{\alpha\gamma}^F(x, x') &= \frac{1}{2} G_{\alpha\gamma}^{(1)}(x, x') + \frac{1}{2} \epsilon(x, x') iD_{\alpha\gamma}(x, x'), \\ &= iG_0^F(x, x') + \frac{1}{2} \left[G_{\alpha\gamma}^{(1)}(x, x') - G_0^{(1)}(x, x') \right],\end{aligned}\tag{4.20}$$

since $D_{\alpha\gamma} = D_0$ can be rewritten in terms of the Feynman propagator. These formulas—and derivations—were first given by Allen [2] for de Sitter. In particular, the Green functions of the two vacua differ only by a homogeneous solution of the wave equation. Alternatively,

$$\begin{aligned}iG_{\alpha\gamma}^F(x, x') &= \cosh^2 \alpha iG_0^F(x, x') + \sinh^2 \alpha iG_0^F(x, x')^* \\ &\quad + \cos \gamma \sinh 2\alpha G_0^{(1)}(x_A, x') - \sin \gamma \sinh 2\alpha D_0(x_A, x').\end{aligned}\tag{4.21}$$

The α -vacuum Feynman propagator is written in terms of Green functions beyond just the Hartle-Hawking Feynman propagator. Also one term involves a complex conjugate, which implies the opposite time ordering.[5, 6] (More precisely, this is the Hartle-Hawking Feynman propagator evaluated at the antipodal points, but with opposite time ordering from that of the antipodal points.)

When $\sin \gamma \neq 0$, the Feynman propagator involves the commutator function. As explained in [2], the commutator function is antisymmetric in x, x' whereas the geodesic distance is symmetric in x, x' . In particular, the commutator function depends on the time ordering and is therefore not invariant under CPT. So the α -vacua are only CPT-invariant for $\gamma = 0, \pi$. [2]

C. Example: The BTZ Black Hole

The three-dimensional BTZ black hole[35, 36] is given by

$$ds^2 = -\left(\frac{r^2}{\ell^2} - M\right)dt^2 + \frac{dr^2}{\frac{r^2}{\ell^2} - M} + r^2 d\phi^2, \quad (4.22)$$

which is the case of $f(r) = \frac{r^2}{\ell^2} - M$; thus $r_+ = \sqrt{M}\Lambda$. In Kruskal coordinates,

$$ds^2 = -4\frac{r_+^2}{M}(1+UV)^{-2}dUdV + r_+^2 \frac{(1-UV)^2}{(1+UV)^2}d\phi^2. \quad (4.23)$$

Spatial infinity is $UV = -1$ and the singularity is at $UV = 1$.

It is well-known that the BTZ black hole is a quotient of AdS_3 . [36] That is, with AdS_3 embedding coordinates X ,

$$-(X^{-1})^2 - (X^0)^2 + (X^1)^2 + (X^2)^2 = -\ell^2, \quad (4.24)$$

we can take

$$\begin{aligned} X^{-1} &= \frac{r}{\sqrt{M}} \cosh \sqrt{M}\phi, & X^0 &= \frac{\ell}{\sqrt{M}} \sqrt{\frac{r^2}{\ell^2} - M} \sinh \frac{\sqrt{M}}{\ell} t, \\ X^1 &= \frac{r}{\sqrt{M}} \sinh \sqrt{M}\phi, & X^2 &= \frac{\ell}{\sqrt{M}} \sqrt{\frac{r^2}{\ell^2} - M} \cosh \frac{\sqrt{M}}{\ell} t, \end{aligned} \quad (4.25)$$

or, for Kruskal coordinates,

$$\begin{aligned} X^{-1} &= \ell \frac{1-UV}{1+UV} \cosh \sqrt{M}\phi, & X^0 &= \ell \frac{U+V}{1+UV} \\ X^1 &= \ell \frac{1-UV}{1+UV} \sinh \sqrt{M}\phi, & X^2 &= \ell \frac{U-V}{1+UV}. \end{aligned} \quad (4.26)$$

With an infinite range for ϕ , these describe AdS_3 ; the BTZ black hole is obtained by identifying $\phi \sim \phi + 2\pi$.

The Green function for a scalar field for the BTZ black hole is obtained by the method of images (see, for example, [37, 38, 39, 40, 41, 42]). For two particles in the same exterior region, with particle 2 on the boundary ($r_2 = \infty$), it reads

$$G(X_1, X_2) = \sum_{n=-\infty}^{\infty} \left[-\sqrt{\frac{r_1^2 - r_+^2}{r_+^2} \cosh\left(\frac{r_+ \Delta t}{\ell^2}\right) + \frac{r_1}{r_+} \cosh \frac{r_+}{\ell} (\Delta\phi + 2\pi n)} \right]^{-2h_+}, \quad (4.27)$$

where $2h_+ = 1 + \sqrt{1 + \ell^2 m^2}$. The authors of [41, 42] noted that this is well-behaved under analytic continuation to X_1 in the other exterior region via $t \rightarrow t - i\frac{\pi\ell^2}{r_+}$; this continues when one additionally takes $\phi \rightarrow \phi + \pi$. That is, the Green function is well-behaved if X_1 is replaced with X_{1A} .

Alternatively, one might note that even pure AdS has a natural antipodal map, defined in the embedding space by

$$(X^{-1}, X^0, X^1, X^2) \longrightarrow (-X^{-1}, -X^0, -X^1, -X^2), \quad (4.28)$$




FIG. 2: The Penrose diagram for AdS_3 , including both $\theta = \pi$ (left-hand side of the diagram) and $\theta = 0$ (the right-hand side). The dashed line is the (spatial) origin of AdS_3 ($\xi = 0$ in Eq. (4.30)). This is AdS_3 , and not its cover. We have drawn two pairs of points that are mapped to each other under the natural AdS_3 antipodal map. This antipodal map correlates timelike-related points, and is not the one we use to construct α -vacua.

and therefore in the BTZ coordinates by

$$t \longrightarrow t - i\frac{\pi\ell^2}{r_+}, \quad \phi \longrightarrow \phi + \frac{\pi}{\sqrt{M}}i. \quad (4.29)$$

Actually, this map is clearer in AdS global coordinates, for which

$$\begin{aligned} X^{-1} &= \ell \cosh \xi \sin \tau, & X^1 &= \ell \sinh \xi \cos \theta, \\ X^0 &= \ell \cosh \xi \cos \tau, & X^2 &= \ell \sinh \xi \sin \theta, \end{aligned} \quad (4.30)$$

and so the map is just

$$\tau \longrightarrow \tau + \pi, \quad \theta \longrightarrow \theta + \pi, \quad (4.31)$$

From the point of view of α -vacua, this is somewhat peculiar; this produces timelike correlations; see Fig. 2. (Also, for AdS, rather than the BTZ black hole, if we are avoiding closed timelike curves by working on the covering space of AdS, this map is isomorphic to $\mathbb{Z}$, not $\mathbb{Z}_2$, and so produces an infinite number of correlations.) More to the point, however, we note that, unlike the antipodal map $(t, \phi) \rightarrow (t - i\frac{\pi\ell^2}{r_+}, \phi + \pi)$, the map (4.29) only changes the Green function (4.27) by a phase, and so does not accomplish anything! Though we have only demonstrated this for the bulk-to-boundary propagator, this is in fact true for the bulk-to-bulk propagator [42, 43, 44]. This is in accord with Witten's demonstration [45] that the pure AdS Green function is essentially unique.

D. The Hartle-Hawking Vacuum

1. Propagators

We now attempt to understand the Feynman propagator in the Hartle-Hawking vacuum, following the original paper [34]. The idea is to look at $G_0^F(x, x')$ on the complexified space-time (T, Z) complex) with x' exterior (in region I) to the black hole and x on the future horizon. General considerations, which we will not review here⁴ imply that, in the complexified spacetime, the propagator is an analytic function of x and x' , except for singularities displaced slightly from null geodesics. Hartle and Hawking write [34]

$$G_0^F(x, x') = K_0(x, x') - i \sum_c \frac{e^{is_c(x, x')/4W_0}}{s_c(x, x') + i\epsilon} D_c(x, x'), \quad (4.32)$$

where c labels the (complex) geodesics connecting x and x' , $s_c(x, x')$ is the geodesic distance along the c th geodesic, K_0 and D_c are analytic functions, W_0 is a small value of the Schwinger parameter, and $\epsilon > 0$. We see that the singularities occur for geodesics that are slightly displaced from null ones.

The question is how x should be perturbed, in the complex plane, in order to sit on the singularity in the propagator (4.32). The general analysis is precisely as in [34]. The singularities of the propagator are in the upper-half V plane and the lower-half U plane. For completeness, we note that the reverse is true for the other exterior region, since the roles of U and V are reversed there.

That the singularities are located in opposite sides of the real axis of the U and V planes implies that the singularities are in the complex t -plane, and not in the complex r -plane. Specifically, the singularities are located to the future of t' and just above the real axis of the complex t -plane, as well as at the image points of these under the $\beta i = \frac{4\pi i}{f'(r_+)}$ periodicity. Moreover since $t + \frac{\beta}{2}i$ corresponds to the antipodal point, the singularity structure is reversed there. In other words, there are also singularities in the past of t' and just below the line $t = \frac{\beta}{2}i$, and its periodic images. This is depicted in Fig. 3(a).

If x is in the same exterior region as x' , then there are real null geodesics for x both in the past of x' as well as the future of x' . Therefore, the singularities are displaced only from the real axis (and its images) as in Fig. 3(b). This is the same singularity structure as for Minkowski space. There are no singularities near $\text{Im } t = \frac{\beta}{2}$, which corresponds to the opposite exterior region and therefore corresponds to points that are spacelike separated from x' .

Now that we know the singularity structure of the Feynman propagator, we can, from (4.11), deduce the singularity structure of the Wightman function (4.8). More precisely, since the Feynman propagator is time ordered, the singularity structure of the Wightman function and the Green function agree for $\text{Re } t > \text{Re } t'$ —at least when this is a sensible time ordering. For the opposite regime of $\text{Re } t$, we can observe that

$$G_0^{(+)}(x, x') = G_0^{(+)}(x', x)^*, \quad (4.33)$$

⁴ Except we note that it is important that the complexified spacetime is nonsingular and complete outside the horizon $r = r_+$. Then the path integral be made well-defined via analytic continuation to the Euclidean regime. We have already checked this. The remaining details appear to be independent of the spacetime in question.




FIG. 3: The singularity structure of $G_0^F(x, x')$ when x' is in region I. For x' in region IV, the figures are shifted vertically by $\frac{\beta}{2}$.




FIG. 4: The singularity structure of $G_0^{(+)}(x, x')$ when x' is in region I. For x' in region IV, the figures are shifted vertically by $\frac{\beta}{2}$.

and so we know the singularity structure. It is depicted in Fig. 4. Note that, unlike the situation for the Feynman propagator, if x' is exterior and x is interior it is impossible to shift t by $\pm\frac{\beta}{2}i$ without crossing a singularity of the Wightman function for some $\text{Re } t$.

2. Thermality

Given this singularity structure of the Green functions, we can examine thermality of the spacetime. The calculation is essentially identical to that in [3]. Explicitly, the transition rate for a detector which follows a trajectory $x'(\tau)$ (τ is proper time) to jump from energy E_i to E_j is given by [46]

$$P_0(E_i \rightarrow E_j) = |m_{ij}|^2 \int_{-\infty}^{\infty} d\tau e^{-i\Delta\epsilon\tau} G_0^{(+)}(x(\tau), x(0)), \quad \Delta\epsilon \equiv \epsilon_j - \epsilon_i, \quad (4.34)$$

where $|m_{ij}|^2$ is a matrix element that depends on the details of the detector. Here $\tau = 0$ has been defined to be the proper time at which $x(0) = x'$, and $\epsilon_{i,j}$ are the proper energies. We consider a detector trajectory at a fixed point exterior to the horizon, $r = R$. Then $\tau = \sqrt{f(R)}(t - t')$ and $\Delta\epsilon = \Delta E / \sqrt{f(R)}$. Since the detector stays in the exterior region, we only need the singularity structure for exterior points, depicted in Fig. 4(b). This singularity structure allows us to shift the contour down to $\text{Im } t = -\frac{\beta}{2}$; thus,

$$P_0(E_i \rightarrow E_j) = |m_{ij}|^2 \sqrt{f(R)} \int_{-\infty}^{\infty} dt e^{-i\Delta E(t-t')} G_0^{(+)}(t, R; t', R), \quad (4.35a)$$

$$= |m_{ij}|^2 \sqrt{f(R)} e^{-\frac{\beta\Delta E}{2}} \int_{-\infty}^{\infty} dt e^{-i\Delta E(t-t')} G_0^{(+)}\left(t - i\frac{\beta}{2}, R; t', R\right). \quad (4.35b)$$

As the point x is in region I, the argument of the Wightman function in (4.35b) corresponds to a point in region IV. Such a point is spacelike separated from the point x' . For spacelike separated points, general principles imply that the commutator function (4.10) vanishes, and the Wightman function is symmetric. Employing time translation invariance—or more precisely, that the Wightman function is a function of the geodesic distance—we can rephrase this statement as

$$G_0^{(+)}\left(t - i\frac{\beta}{2}, R; t', R\right) = G_0^{(+)}\left(t' - t + i\frac{\beta}{2}, R; 0, R\right), \quad (4.36)$$

since both sides of the equation are a function of the same geodesic distance, and there are no nearby singularities. Upon using eq. (4.36), βi periodicity and replacing the integration variable t by $t' - t$, equation (4.35b) reads

$$\begin{aligned} P_0(E_i \rightarrow E_j) &= |m_{ij}|^2 \sqrt{f(R)} e^{-\frac{\beta\Delta E}{2}} \int_{-\infty}^{\infty} dt e^{i\Delta E t} G_0^{(+)}\left(t - i\frac{\beta}{2}, R; 0, R\right), \\ &= |m_{ij}|^2 \sqrt{f(R)} e^{-\beta\Delta E} \int_{-\infty}^{\infty} dt e^{i\Delta E t} G_0^{(+)}(t, R; 0, R), \end{aligned} \quad (4.37)$$

upon shifting the contour again. In other words, using (4.35a) on the right-hand side,

$$\frac{P_0(E_i \rightarrow E_j)}{P_0(E_j \rightarrow E_i)} = e^{-\beta\Delta E}. \quad (4.38)$$

This is the condition for detailed balance at temperature

$$T = \beta^{-1} \quad (4.39)$$

as predicted from the periodicity of Euclidean time.

E. Thermality of α -Vacua

Let us repeat the calculation of detailed balance for an α -vacuum. We again start with

$$P_{\alpha\gamma}(E_i \rightarrow E_j) = |m_{ij}|^2 \sqrt{f(R)} \int_{-\infty}^{\infty} dt e^{-i\Delta E(t-t')} G_{\alpha\gamma}^{(+)}(t, R; t', R); \quad (4.40)$$

the difference between (4.34) and (4.40) is in the Wightman function. Substituting (4.16), we find

$$\begin{aligned} P_{\alpha\gamma}(E_i \rightarrow E_j) = |m_{ij}|^2 \sqrt{f(R)} \int_{-\infty}^{\infty} dt e^{-i\Delta E(t-t')} \Big\{ & \cosh^2 \alpha G_0^{(+)}(t, R; t', R) \\ & + \frac{1}{2} e^{i\gamma} \sinh 2\alpha G_0^{(+)}\left(t - i\frac{\beta}{2}, R; t', R\right) + \frac{1}{2} e^{-i\gamma} \sinh 2\alpha G_0^{(+)}\left(t, R; t' + i\frac{\beta}{2}, R\right) \\ & + \sinh^2 \alpha G_0^{(+)}\left(t - i\frac{\beta}{2}, R; t' + i\frac{\beta}{2}, R\right) \Big\} \end{aligned} \quad (4.41)$$

Again, since for spacelike separated points, the Wightman function depends only on the geodesic distance, we can use

$$G_0^{(+)}\left(t, R; t' + i\frac{\beta}{2}, R\right) = G_0^{(+)}\left(t - i\frac{\beta}{2}, R; t', R\right), \quad (4.42)$$

for the middle two terms, and appropriately shift the contour. For the last term, we shift the contour⁵ and then use (4.42) and shift the contour again. As a result,

$$P_{\alpha\gamma}(E_i \rightarrow E_j) = |m_{ij}|^2 \sqrt{f(R)} \left| \cosh \alpha + \sinh \alpha e^{i\gamma} e^{\frac{\beta}{2}\Delta E} \right|^2 P_0(E_i \rightarrow E_j). \quad (4.43)$$

Thus,

$$\frac{P_{\alpha\gamma}(E_i \rightarrow E_j)}{P_{\alpha\gamma}(E_j \rightarrow E_i)} = \left| \frac{\cosh \alpha + \sinh \alpha e^{i\gamma} e^{\frac{\beta}{2}\Delta E}}{\cosh \alpha + \sinh \alpha e^{i\gamma} e^{-\frac{\beta}{2}\Delta E}} \right|^2 e^{-\beta\Delta E}. \quad (4.44)$$

This is not thermal. After taking into account the conventions described in footnote 3 (p. 11), it matches the nonthermal expression in [3]. Indeed, the authors of [4] have already observed that the de Sitter result depends only on the analytic structure of the Green function.

V. (NO) OBJECTIONS TO α -VACUA

There has been a substantial amount of work attempting to debunk the idea of α -vacua for de Sitter space. In this section we will address the objections and describe how they may be evaded in black hole spacetimes.

We will see that some of the objections involve consideration of spacetime points that are separated by the black hole horizon. Issues involving causality and correlation functions of

⁵ Here x' is in region IV, so the poles in t will also be in region IV; *i.e.* just off the $\text{Im } t = -\beta i/2$ line, and its images. Moreover, time is reversed in region IV, so the poles are just below this line—the opposite of Fig. 4(b) in which the poles are just above $\text{Im } t = 0$ and its images. Therefore we can shift the t contour up to the real axis without crossing any poles.




FIG. 5: Perturbations of Schwarzschild black holes increase the area of the horizon. On the Penrose diagram, this translates to the depicted deformation of the future singularity. This particular diagram is the “eternal Vaidya” black hole that is obtained by taking an eternal Schwarzschild black hole of mass M and collapsing a spherical shell of radially directed null matter of additional energy M , just to the future of the past horizon $v = -\infty$.

such points are already poorly understood, and are related to questions involving information loss in, and the end-point of, Hawking radiation. For example [47, 48], the concept of black hole complementarity has been suggested as a means to try to assert the consistency of black hole physics. According to black hole complementarity, one may describe a black hole system as seen by an observer who falls through the horizon and sees no irregularities as she falls through the horizon, or as seen by an observer constrained to live outside the horizon and who sees the Hawking radiation and other thermal physics of the black hole, but one may not describe the black hole using both observers as these are complementary descriptions. This avoids, for example, the quantum Xerox problem (*e.g.* [49])—that black hole evaporation combined with the causal disconnect between the two sides of the black hole horizon requires that the quantum information of the infalling matter be copied to outside states, in conflict with the unitarity of quantum mechanics—but also leaves open the question of how one should treat quantities such as correlation functions between points inside and outside the black hole horizon.

Thus, since physics connecting the interior and exterior of black holes is poorly understood, even for the well-known vacua, we should not expect to have resolutions for puzzles which arise in such a context for α -vacua. As a result, we will be brief in discounting such problems in the following.

A. Causality

Naïvely, because an exterior point x and its antipodal point x_A are spacelike separated, no causality problems arise from correlating the two points. More precisely, for de Sitter spaces, there are naïvely no causality problems because the lightcones emanating from x and x_A do not intersect. However, it has been noted [5] that this does not take backreaction into account. Backreaction acts to increase the de Sitter horizon and therefore makes de Sitter “taller”. [50] Then the lightcones emanating from x and x_A do intersect in the future, and this leads to causal problems for α -vacua.

For black holes spacetimes, however, the backreaction will increase the black hole horizon, thus increasing the amount of the space behind the black hole horizons. That is, the future

singularity in Fig. 1 arcs upwards (see Fig. 5). Thus, the causality problems that appear due to intersection of the lightcones emanating from x in region I and x_A in region IV only occur inside the black hole horizon. Since we already do not understand physics inside the horizon, this does not bother us.

B. Poles

In field theory, there are singularities in tree-level three-point functions, $\langle \phi(x_1)\phi(x_2)\phi(x_3) \rangle$. Namely, since the two-point functions are singular for null separations (*cf.* Eq. (4.32)), and since the three-point function involves an integral,

$$\langle \phi(x_1)\phi(x_2)\phi(x_3) \rangle \sim \int dy G^F(x_1, y) G^F(x_2, y) G^F(x_3, y), \quad (5.1)$$

there are poles when the integrated interaction point y is on the lightcone of one of the external points. Of course, these poles are well-understood in terms of physical propagating particles. However, for α -vacua, such poles also appear when y is on the lightcone of the antipodal of one of the external points.[5] This appears to be unphysically acausal—in principle, it allows an observer to probe a causally disconnected region, namely that of the antipodal points, if all the points x_1, x_2, x_3 are causally connected. Moreover, if one of the points happens to also lie on the lightcone of another's the antipodal point, then for y also on that lightcone, there will be coincident poles leading to divergence of the tree-level diagram. These issues lead to questions about the sensibleness of α -vacua.

For black hole spacetimes, however, one again sees from Fig. 1 that this problem only arises if at least one of x_1, x_2, x_3 is inside the black hole horizon, as this is the only place where the causal future of the exterior regions intersects. Since we already do not understand correlation functions between fields inside and outside the horizon, this does not bother us.

C. Thermality

The nonthermality of Eq. (4.44) appears to be a disaster [4]. Assuming a steady state system, transitions from the i th state to the j th state should be offset by transitions from the j th state to the i th. This is the statement of detailed balance:

$$\rho(E_i)P(E_i \rightarrow E_j) = \rho(E_j)P(E_j \rightarrow E_i) \Leftrightarrow \frac{P(E_i \rightarrow E_j)}{P(E_j \rightarrow E_i)} = \frac{\rho(E_j)}{\rho(E_i)}. \quad (5.2)$$

For the $\alpha = 0$ vacuum, we conclude $\rho(E_i) \propto e^{-\beta E_i}$, where β is the inverse temperature, in appropriate units. This is just the Boltzmann distribution.

Indeed, as the authors of [4] have reminded us, detailed balance (5.2) implies the Boltzmann distribution. Setting

$$R_{ij} \equiv \frac{P(E_i \rightarrow E_j)}{P(E_j \rightarrow E_i)} \equiv \varphi(E_i - E_j), \quad (5.3)$$

detailed balance implies that

$$R_{ij}R_{jk} = R_{ik}, \quad (5.4)$$

so if $\delta E \equiv E_j - E_i = E_k - E_j$ then

$$\varphi(\delta E)^2 = \varphi(2\delta E), \quad (5.5)$$

which implies $\varphi(\delta E) = e^{\varphi'(0)\delta E}$. That is just a Boltzmann distribution, upon identifying $\beta = -\varphi'(0)$.

However, there are loopholes. For one, we know that the Boltzmann distribution is the classical distribution. For Fermi-Dirac statistics, for example, the transition from the i th state to the j th state cannot occur if the j th state is already occupied. That is, Eq. (5.2) is replaced by

$$\rho(E_i) [1 - \rho(E_j)] P(E_i \rightarrow E_j) = \rho(E_j) [1 - \rho(E_i)] P(E_j \rightarrow E_i) \quad (5.6)$$

or

$$\frac{P(E_i \rightarrow E_j)}{P(E_j \rightarrow E_i)} = \frac{\rho(E_j) [1 - \rho(E_i)]}{\rho(E_i) [1 - \rho(E_j)]}. \quad (5.7)$$

Indeed, applying this to the $\alpha = 0$ result (4.38) gives

$$\rho(E_i) \propto \frac{e^{-\beta E_i}}{1 + e^{-\beta E_i}}, \quad (5.8)$$

which is the familiar Fermi-Dirac distribution. The Bose distribution is similarly obtained by realizing that the transition for bosons is enhanced if the excited state is already occupied.[51]

Nevertheless, it is clear that for a general α -vacuum, the result (4.44) will not correspond to a familiar distribution. For one, the expression does not factorize into a product of functions of E_i and E_j ! This is also not a disaster, however. Most conservatively, we need not presume that the distribution in question is that of an equilibrium system. It is well-known (see *e.g.* [52]) that the late time limit of any nonequilibrium distribution function is a time-independent nonequilibrium steady-state distribution function. This need not be a familiar distribution; it need only solve the Fokker-Planck equation (or the appropriate quantum generalization thereof) which, being a second-order differential equation, has a time-independent solution that is not the Boltzmann distribution. That is apparently the situation here. A concrete example of such a situation appears in astrophysics [53].

An additional motivation for realizing that the result (4.44) corresponds to a nonequilibrium steady-state system is the following. Consider the instanton that is the Euclidean black hole. Naïvely, its temperature is determined by a length scale, namely the radius of the instanton. For the α -vacua, modes are explicitly correlated between the two “sides” of the space time, permitting a “short cut” across the instanton. Presumably, this affects the thermality.

However, we cannot resist mentioning that one can obtain a familiar thermal distribution by replacing the detailed balance relation (5.2) with the peculiar expression⁶

$$\begin{aligned} & \left| \cosh \alpha \sqrt{\rho(E_i) [1 - \sigma \rho(E_j)]} + e^{i\gamma} \sinh \alpha \sqrt{\rho(E_j) [1 - \sigma \rho(E_i)]} \right|^2 P(E_i \rightarrow E_j) \\ &= \left| \cosh \alpha \sqrt{\rho(E_j) [1 - \sigma \rho(E_i)]} + e^{i\gamma} \sinh \alpha \sqrt{\rho(E_i) [1 - \sigma \rho(E_j)]} \right|^2 P(E_j \rightarrow E_i), \end{aligned} \quad (5.9)$$

⁶ This is clearly not unique.

which implies the distribution

$$\rho(E_i) \propto \frac{1}{e^{\beta E_i} - \sigma}. \quad (5.10)$$

This is a Boltzmann distribution for $\sigma = 0$, a Bose-Einstein distribution for $\sigma = 1$ and a Fermi-Dirac distribution for $\sigma = -1$, all at temperature β . Eq. (5.9) would imply that the meaning of detailed balance—*i.e.* the quantum statistics—is vacuum dependent, as for anyons. However, there is still at least one puzzle which follows from this idea—namely, for $\alpha \neq 0$ it would be possible to excite the detector from state $|i\rangle$ to $|j\rangle$ even if the initial population of the $|i\rangle$ 'th state was empty provided $|j\rangle$ was sufficiently populated—and though that puzzle may have a resolution in terms of an interpretation of a transition from energy E_j to E_i at the antipodal point, we will not advocate this interpretation as we prefer the nonequilibrium statistical mechanical interpretation of (4.44).

D. Pinch Singularities

It has been noted [5]—see [9] for a more general phrasing of this result—that quantum field theory loops become ill-defined for α -vacua because of the emergence of pinch singularities. For example, consider the one-loop correction to the propagator

$$\begin{aligned} \text{---} x \text{---} \bigcirc \text{---} y \text{---} &\sim \int dx \int dy G_{\alpha\gamma}^F(x, y) G_{\alpha\gamma}^F(y, x) \\ &\sim \dots + \int dx \int dy \cosh^2 \alpha \sinh^2 \alpha G_0^F(x, y) G_0^F(y, x)^* + \dots \end{aligned} \quad (5.11)$$

The term we have shown involves both $i\epsilon$ prescriptions, and thus it is impossible to deform the contour of integration so as to avoid the singularities. More precisely, the residue of the pole from any one factor is singular due to the pole in the other term.[5]

For this reason, many authors [6, 7, 8, 10] have suggested modifying the time ordering prescription for α -vacua, in order to remove the pinch singularities. But presumably string quantization works in spacetimes with nonpositive cosmological constant, and it is well-known that string loops are nonsingular. Thus, we anticipate that for the black hole spacetimes considered here, the pinch singularities can be evaded without altering the time ordering prescription.

That this anticipation is not ridiculous is seen by considering strings in Rindler space.⁷ Rindler space is just Minkowski space after a coordinate transformation, and string n -point amplitudes are just integrated worldsheet correlation functions,

$$\langle \mathcal{V}_1 \cdots \mathcal{V}_n \rangle. \quad (5.12)$$

⁷ Past work on strings in Rindler space includes [54, 55] but we have not seen the approach given here, which resembles some calculations of [56], included in the Rindler literature.




FIG. 6: The (a) planar and (b) nonplanar 1-loop open string 2-point diagrams. As *position space field theory* diagrams (take widths to zero) the two internal propagators connecting the two vertices, when expanded in an α -vacuum, have cross-terms that give the pinch singularities. However, pulling out the right-hand external state for the nonplanar diagram (b) gives a single closed string propagator, and therefore no pinch singularity. So any pinch singularity must come from the opposite limit, which is the field theory limit, $t \rightarrow \infty$.

For tachyons in ordinary Minkowski space, the vertex operators are just the Minkowski modes $\mathcal{V}_j = e^{ik_j \cdot X}$; in Rindler space, they are the Rindler modes. The latter, however, are obtainable as a Bogliubov transformation of the former; that is, string amplitudes in Rindler space can be obtained as an appropriate integral (over $k_1, \dots, k_n$) of the Minkowski string amplitudes. This yields a finite result. One can define “ α -vacua” of Rindler space using PT as the “antipodal map”, though they are not as well-defined because PT has a fixed point at the origin of the Minkowski spacetime. Although loop amplitudes in these “ α -vacua” would naïvely be expected to suffer pinch singularities due to the reversal of Rindler time, in fact the string loop amplitudes are just linear combinations of the ordinary, and finite, Rindler space amplitudes, and so are finite. We conclude that strings do not suffer from pinch singularities. The still-skeptical reader can find details in Appendix B.

One might still ask⁸ how it is that the field theory pinch singularities are recovered from the finite string theory amplitudes. For concreteness, we will consider the open string in our Rindler toy model. The open string one-loop annulus diagram (*e.g.* Fig. 6) is conformally equivalent to a cylinder, and includes an integral over the modulus t , the ratio between the radius and the length of the cylinder. The field theory limit is obtained in the limit $t \rightarrow \infty$, in which the open string loop becomes infinitely short. The annulus diagram also includes integrals over the positions of the vertex operators which live on the boundary; these range over the circumference of the cylinder, $2\pi t$.

By considering open string factorization of the annulus diagram, (this argument follows [57] most closely and is reproduced in Appendix B 3 b i, although this result dates back to at least [58])—namely, that the one-loop annulus amplitude can be reconstructed from the tree-level amplitude on the plane by insertion of two open string vertices in the plane, as depicted in Fig. 7—one can recover the propagator of the intermediate particle. The additional propagator factor which leads to the pinch singularity is obtained from the integral over the vertex operator positions as well. However, the pinch singularity is only obtained because in the $t \rightarrow \infty$ limit, the vertex operator position extends, in the coordinates of Appendix B 3 b i, to $y = \pm 1$. For finite t , the region of integration for the vertex operator position integral is shrunk to $y = \pm \tanh \frac{\pi t}{2}$, and no singularity appears. In this way, string

⁸ J.M thanks the referee for bringing this omission in our original argument to our attention.




FIG. 7: Open string factorization of the annulus occurs as the modular parameter of the annulus $t \rightarrow \infty$, corresponding to an infinitely short cylinder. For a general value of the modulus, the annulus diagram is equivalent to a sum of tree-level diagrams in which one inserts vertex operators with equal and opposite momentum and cuts out, and identifies, a region around the inserted vertex operators. The sum is over the momentum and identity of the inserted vertex operators; this corresponds to the loop exchange of a virtual string. This figure was stolen from [57].

theory regularizes the field theory pinch singularity.

On this note, it is intriguing to recall (see *e.g.* [59, 60, 61]) that naïvely expected pinch singularities in nonequilibrium field theory fail to appear after careful calculation if the interaction that leads to the pinch singularity is turned on for only a finite time. Since the integration over the location of the string theory vertex operator is an integration over a worldsheet time coordinate, the finiteness of the modulus t , away from the field theory limit, corresponds to finiteness of the worldsheet time and so could be precisely this mechanism. We hope that further work will shed light on this.

E. Uniqueness of Black Hole Vacua

It is known [62] that of the usually considered black hole vacua, only the Hartle-Hawking vacuum enjoys the property that the renormalized stress tensor does not diverge on either the past or future black hole horizon. Does this not rule out α -vacua?⁹

Actually, the stress tensor does not diverge on the horizon in any α -vacuum constructed from the Hartle-Hawking vacuum.¹⁰ The point is that the renormalization of the stress tensor is related to $G_{\alpha\gamma}^F(x, x)$, where x is evaluated on the black hole horizon. Using (4.21), this will be finite if the Hartle-Hawking Green functions $G_0^F(x, x)$, $G_0^{(1)}(x, x_A)$ and $D_0(x, x_A)$ are finite. The work of Candelas [62] demonstrated that $G_0^F(x, x)$ is finite on the horizon. Moreover, for x on the black (white) hole horizon, x_A is on the white (black) hole horizon; on the Penrose diagram, the points appear to be null-like related, but because the antipodal map includes the antipodal map on the sphere, x and x_A are therefore actually spacelike related for x on the horizon. Thus $D_0(x, x_A) = 0$, and $G_0^{(1)}(x, x_A) = G_0^F(x, x_A)$ is finite. So the renormalized stress tensor is finite on the horizon for all α -vacua constructed from the Hartle-Hawking vacuum.

This does not contradict Candelas' result, as he only considered the Boulware, Hartle-Hawking and Unruh vacua.

⁹ J.M. thanks Rob Myers for asking this interesting question at the Andrew Chamblin Memorial Symposium at the University of Louisville in Kentucky, USA.

¹⁰ Similarly, the stress tensor diverges on the past but not the future horizon in any α -vacuum constructed from the Unruh vacuum.

VI. CFT α -STATES

Given that Schwarzschild-AdS has two boundaries and α -vacua, there should be a one-parameter set of CFT states. We provide a proposal in this section.

First, though, we should point out that this proposal differs substantially from that for dS/CFT [3]. For dS/CFT, the authors noted that, unlike AdS/CFT¹¹, each dS mode is associated with two CFT operators, the product of which is marginal. Thus, the dS α -vacua can be associated with the CFT deformed by this marginal operator. For AdS, however, because only one of the two candidate CFT operators actually exists¹¹, this prescription does not apply.

So let us recall the prescription [26, 32] for the $\alpha = 0$ vacuum. The two boundary CFTs give a product Hilbert space $|\rangle_1|\rangle_2$. Because time runs in opposite directions on the two boundaries, one has, just as in real-time thermal field theory (for a review see [64]), $H = H_1 \otimes \mathbb{1} - \mathbb{1} \otimes H_2$. Thus the state

$$|\psi\rangle_0 = \frac{1}{\sqrt{Z}} \sum_i e^{-\beta E_i/2} |i\rangle_1 |i\rangle_2, \quad (6.1)$$

written in terms of a complete set of energy eigenstates of each CFT, has vanishing energy. It is unit normalized using the partition function Z . It cannot be overemphasized that it is a *pure state*.

The two identical boundary CFTs are related by an antilinear involutive map [64], with the result that

$$e^{\beta H/2} [\mathbb{1} \otimes \mathcal{O}(t, \Omega)] |\psi\rangle_0 = [\mathcal{O}^\dagger(t, \Omega) \otimes \mathbb{1}] |\psi\rangle_0, \quad (6.2a)$$

or equivalently, and somewhat more schematically,

$$\mathcal{O}_2(t, \Omega) |\psi\rangle_0 = \mathcal{O}_1^\dagger(t - i\frac{\beta}{2}) |\psi\rangle_0, \quad (6.2b)$$

Here, $\mathcal{O}_1^\dagger(t - i\frac{\beta}{2}) = [e^{H_1\beta/2} \mathcal{O}_1(t) e^{-H_1\beta/2}]^\dagger$ is the Hermitian conjugate of the operator evaluated at time $t - i\frac{\beta}{2}$ and so is the Hermitian conjugate of the operator, evaluated at time $t + i\frac{\beta}{2}$. As a result, one has the relation, for operators A, B in each field theory, (*cf.* [32])

$$\langle \psi |_0 A_1(t) B_2(t') | \psi \rangle_0 = \langle \psi |_0 A_1(t) B_1^\dagger(t' - i\frac{\beta}{2}) | \psi \rangle_0. \quad (6.3)$$

However, observers on one boundary or the other can only see their own boundary. Such an observer, in CFT₁ for definiteness, will not see boundary 2, and so its observations will involve a trace over the CFT₂ Hilbert space. That is, an observer in CFT₁ sees the density matrix

$$\rho_1 = \text{Tr}_2 |\psi\rangle \langle \psi| = \frac{1}{Z} \sum_i e^{-\beta E_i} |i\rangle_1 {}_1\langle i|. \quad (6.4)$$

¹¹ The exception is the special range, $-\frac{d^2}{4} < m^2 < -\frac{d^2}{4} + 1$, of AdS_{d+1} tachyon masses for which the two CFT operators, $\mathcal{O}_\pm$, correspond to two inequivalent bulk quantizations of the scalar field.[63] In particular, it is argued in [63] that the two inequivalent quantizations of AdS correspond to two different CFTs, one with only $\mathcal{O}_+$ and one with only $\mathcal{O}_-$. So even in this special range, the putatively marginal $\mathcal{O}_+\mathcal{O}_-$ does not exist in AdS. And even if this operator did exist in this case, it would be unsatisfying to have a CFT α -state prescription that only held in this limited range of AdS masses.

This is precisely the thermal density matrix.

The Schwarzschild-AdS α -vacua are constructed by a Bogoliubov transformation that correlates antipodal points. In the CFT, then, this suggests an analogous Bogoliubov transformation. We will make this precise for a single one-dimensional, harmonic oscillator on each “boundary”; the generalization to actual dual CFTs should be obvious by replacing the harmonic oscillator creation and annihilation operators by operators which create and annihilate the state $|i\rangle$ on each boundary.

We have two copies of the harmonic oscillator,

$$[a_1, a_1^\dagger] = 1 = [a_2, a_2^\dagger]. \quad (6.5)$$

The pure state above can be written as

$$|\psi\rangle_0 = \sqrt{1 - e^{-\beta}} \exp \left[e^{-\frac{\beta}{2}} a_1^\dagger a_2^\dagger \right] |0\rangle_1 |0\rangle_2, \quad (6.6)$$

after setting the harmonic oscillator frequency $\omega = 1$ (otherwise $\beta \rightarrow \beta\omega$) and using

$$Z = \frac{1}{2} \operatorname{csch} \frac{\beta}{2}. \quad (6.7)$$

We can now perform the Bogoliubov transformation,

$$b_1^\dagger = \cosh \alpha a_1^\dagger - e^{i\gamma} \sinh \alpha a_2, \quad b_2^\dagger = \cosh \alpha a_2^\dagger - e^{i\gamma} \sinh \alpha a_1. \quad (6.8)$$

Note that this preserves the Hamiltonian

$$H = a_1^\dagger a_1 - a_2^\dagger a_2 = b_1^\dagger b_1 - b_2^\dagger b_2, \quad (6.9)$$

in accord with the statement that AdS α -vacua preserve the symmetries preserved by the standard (Hartle-Hawking) vacuum. The new operators define new vacua via

$$b_1 |\tilde{0}\rangle_1 = 0 = b_2 |\tilde{0}\rangle_2. \quad (6.10)$$

Note that although the “CFT” is still a product CFT, it is no longer manifestly a product of CFTs on each boundary.

We now use the Bogoliubov transformed operators to build a new pure state, $|\psi\rangle_{\alpha\gamma}$, namely

$$|\psi\rangle_{\alpha\gamma} = \sqrt{1 - e^{-\beta}} \exp \left[e^{-\frac{\beta}{2}} b_1^\dagger b_2^\dagger \right] |\tilde{0}\rangle_1 |\tilde{0}\rangle_2. \quad (6.11)$$

To what density matrix does this state correspond? Again, we consider an observer on the first boundary. Such an observer does not observe the second boundary, and so we should again trace over CFT_2 . This is *not* the same as tracing over the $(b_2^\dagger, b_2)$ Hilbert space! That Hilbert space partly exists on CFT_1 . That is,

$$\operatorname{Tr}_2 \neq \widetilde{\operatorname{Tr}}_2. \quad (6.12)$$

In Appendix A we show that

$$|\psi\rangle_{\alpha\gamma} = \frac{\operatorname{sech} \alpha \sqrt{1 - e^{-\beta}}}{1 + e^{-\frac{\beta}{2} + i\gamma} \tanh \alpha} \exp \left[\frac{e^{-\frac{\beta}{2}} + e^{-i\gamma} \tanh \alpha}{1 + e^{-\frac{\beta}{2} + i\gamma} \tanh \alpha} a_1^\dagger a_2^\dagger \right] |0\rangle_1 |0\rangle_2. \quad (6.13)$$




FIG. 8: The α dependence of the entropy at extreme (a) low temperature and (b) high temperature.

We should note, following [6], that this makes the state associated with α -vacua appear to be in the same Hilbert space as the state associated with the Hartle-Hawking vacuum. However, the CFT associated with an AdS black hole has an infinite number of oscillators; then the inner product ${}_{\alpha\gamma}\langle\psi|\psi\rangle_0 \sim \text{sech}^\infty \alpha = 0$. This argument is easily generalized to any state built up from $|\psi\rangle_{\alpha\gamma}$ and so states for different values of α, γ are in different Hilbert spaces.

Eq. (6.13) yields the density matrix

$$\begin{aligned} \rho_{\alpha\gamma} &= \text{Tr}_2 |\psi\rangle_{\alpha\gamma} {}_{\alpha\gamma}\langle\psi| \\ &= (1 - e^{-\beta}) \frac{\text{sech}^2 \alpha}{\left|1 + e^{-\frac{\beta}{2} + i\gamma} \tanh \alpha\right|^2} \sum_{n=0}^{\infty} e^{-\beta n} \left| \frac{\cosh \alpha + e^{\frac{\beta}{2} + i\gamma} \sinh \alpha}{\cosh \alpha + e^{-\frac{\beta}{2} + i\gamma} \sinh \alpha} \right|^{2n} |n\rangle_1 {}_1\langle n| \end{aligned} \quad (6.14)$$

Note that at $\alpha = 0$, this reduces to the standard thermal answer. Also, this shows that the density matrix is α -dependent, and thus the entropy $S_{\alpha\gamma} = -\text{Tr} \rho_{\alpha\gamma} \ln \rho_{\alpha\gamma}$ is α -dependent.

Explicitly,

$$S_{\alpha\gamma} = -\ln \left[\frac{(1 - e^{-\beta}) \text{sech}^2 \alpha}{\left|1 + e^{-\frac{\beta}{2} - i\gamma} \tanh \alpha\right|^2} \right] - \frac{\left|e^{-\frac{\beta}{2}} + e^{i\gamma} \tanh \alpha\right|^2 \cosh^2 \alpha}{1 - e^{-\beta}} \ln \frac{\left|e^{-\frac{\beta}{2}} + e^{i\gamma} \tanh \alpha\right|^2}{\left|1 + e^{-\frac{\beta}{2} - i\gamma} \tanh \alpha\right|^2}. \quad (6.15)$$

It is interesting to see how α affects the high and low temperature entropy. This is shown in Fig. 8.

VII. ADS/CFT

A. Populations

Let us compare Eq. (6.14) and (4.44). To do this, we should make the usual identification between the harmonic oscillator occupation number, and the number of particles in a

given state. Thus, [upon restoring the frequency to Eq. (6.14) via $\beta \rightarrow \beta E_i$] the ratio of probabilities between having n particles in the i^{th} state and $n - 1$ particles in that CFT state, is

$$\frac{\rho(n)}{\rho(n-1)} = e^{-\beta E_i} \left| \frac{\cosh \alpha + e^{\frac{\beta E_i}{2} + i\gamma} \sinh \alpha}{\cosh \alpha + e^{-\frac{\beta E_i}{2} + i\gamma} \sinh \alpha} \right|^2, \quad (7.1)$$

which exactly matches the ratio of transition rates of the bulk detector (4.44)!

B. Propagators

Now let us make the prescription of §VI somewhat more precise. In [65, 66] it is argued that the mode expansion of the bulk field ϕ ,

$$\phi(x) = \sum_{\kappa, n, s(n)} \left[a_{\kappa, n, s(n)} \phi_{\kappa, n, s(n)}(x) + a_{\kappa, n, s(n)}^\dagger \phi_{\kappa, n, s(n)}^*(x) \right], \quad (7.2)$$

implies the mode expansion of the dual operator

$$\mathcal{O}(t, \Omega) = \sum_{\kappa, n, s(n)} \left[b_{\kappa, n, s(n)} \tilde{\phi}_{\kappa, n, s(n)}(t, \Omega) + b_{\kappa, n, s(n)}^\dagger \tilde{\phi}_{\kappa, n, s(n)}^*(t, \Omega) \right], \quad (7.3)$$

where the mode $\tilde{\phi}_{\kappa, n, s(n)}$ is related to the boundary value of $\phi_{\kappa, n, s(n)}$ and Ω parametrizes the boundary S^{d-1} . With two boundaries, this is more subtle. We would like to write, on each boundary,

$$\begin{aligned} \mathcal{O}_1(t, \Omega) &= \sum_{\kappa, n, s(n)} \left[b_{1\kappa, n, s(n)} \tilde{\phi}_{\kappa, n, s(n)}(t, \Omega) + b_{1\kappa, n, s(n)}^\dagger \tilde{\phi}_{\kappa, n, s(n)}^*(t, \Omega) \right], \\ \mathcal{O}_2(t, \Omega) &= \sum_{\kappa, n, s(n)} \left[b_{2\kappa, n, s(n)} \tilde{\phi}_{\kappa, n, s(n)}(t - i\frac{\beta}{2}, \Omega) + b_{2\kappa, n, s(n)}^\dagger \tilde{\phi}_{\kappa, n, s(n)}^*(t - i\frac{\beta}{2}, \Omega) \right], \end{aligned} \quad (7.4)$$

but the bulk modes, restricted to each boundary, form an overcomplete set of functions on each boundary. Basically, the bulk modes come in pairs, one linear combination of which vanishes on one boundary and another of which vanishes on the other boundary. However, we can write Eq. (7.4) assuming that we have restricted the sums to a complete set of functions, and such that, just as for the bulk theory, we have

$$\tilde{\phi}_{\kappa, n, s(n)}^*(t, \Omega) = \tilde{\phi}_{\kappa, n, s(n)}(t - i\frac{\beta}{2}, \Omega_A), \quad (7.5)$$

where Ω_A is the antipodal point on the sphere to Ω , and $t - i\frac{\beta}{2}$ is the corresponding time on the other boundary. Moreover, [67] (see also [42, 65, 66, 68]) argue that the quantum numbers that we have incorporated into κ are essentially the energy (after all, there is a time-like Killing vector) and that the relationship Eqs. (7.2) and (7.3) implies that the two-point function $\langle \mathcal{O}(t, \Omega) \mathcal{O}(t', \Omega') \rangle$ is given by the bulk propagator. We would now like to check that this holds as well after the Bogoliubov transformation.

First, however, we should note that since the operators in the mode expansion of the composite operator (7.3) have well-defined energies, that the compositeness of the operator to which the creation and annihilation operators are defined is not incompatible with our

previous discussions in §VI, §VII A. Indeed, it will be sufficient to discuss a single set of oscillators, so that

$$\begin{aligned}
\mathcal{O}_1 &= A_1(t, \Omega) a_1 + A_1(t, \Omega)^* a_1^\dagger \\
&= A_1 \left(\cosh \alpha b_1 + e^{i\gamma} \sinh \alpha b_2^\dagger \right) + A_1^* \left(\cosh \alpha b_1^\dagger + e^{-i\gamma} \sinh \alpha b_2 \right), \\
\mathcal{O}_2 &= A_2(t - i\frac{\beta}{2}, \Omega) a_2 + A_2(t - i\frac{\beta}{2}, \Omega)^* a_2^\dagger \\
&= A_2 \left(\cosh \alpha b_2 + e^{i\gamma} \sinh \alpha b_1^\dagger \right) + A_2^* \left(\cosh \alpha b_2^\dagger + e^{-i\gamma} \sinh \alpha b_1 \right).
\end{aligned} \tag{7.6}$$

Let us note that

$${}_{\alpha\gamma} \langle \psi | b_1 b_1^\dagger | \psi \rangle_{\alpha\gamma} = \frac{1}{2} e^{\frac{\beta}{2}} \operatorname{csch} \frac{\beta}{2}, \quad {}_{\alpha\gamma} \langle \psi | b_1^\dagger b_1 | \psi \rangle_{\alpha\gamma} = \frac{1}{2} e^{-\frac{\beta}{2}} \operatorname{csch} \frac{\beta}{2}, \quad {}_{\alpha\gamma} \langle \psi | b_1 b_2 | \psi \rangle_{\alpha\gamma} = \frac{1}{2} \operatorname{csch} \frac{\beta}{2}, \tag{7.7}$$

Therefore, for $t > t'$,

$$\begin{aligned}
{}_{\alpha\gamma} \langle \psi | \mathcal{O}_1(t, \Omega) \mathcal{O}_1(t', \Omega') | \psi \rangle_{\alpha\gamma} &= \cosh^2 \alpha {}_0 \langle \psi | \mathcal{O}_1(t, \Omega) \mathcal{O}_1(t', \Omega') | \psi \rangle_0 \\
&+ \sinh^2 \alpha {}_0 \langle \psi | \mathcal{O}_1(t, \Omega) \mathcal{O}_1(t', \Omega') | \psi \rangle_0^* + \cos \gamma \sinh 2\alpha {}_0 \langle \psi | \mathcal{O}_1(t, \Omega) \mathcal{O}_1(t' - i\frac{\beta}{2}, \Omega'_A) | \psi \rangle_0.
\end{aligned} \tag{7.8}$$

The form of the last term arises by identifying $A_1(t, \Omega) = A_1(t - i\frac{\beta}{2}, \Omega_A)^*$ and by noting that the correlation function between the point (t, Ω) and the “antipodal” point $(t' - i\frac{\beta}{2}, \Omega'_A)$ should be identified with ${}_0 \langle \psi | \mathcal{O}_1(t, \Omega) \mathcal{O}_2(t', \Omega') | \psi \rangle_0$; in particular, since this corresponds to the bulk propagator between spacelike points, it should be identified with the Hadamard function. Thus, this agrees precisely with Eq. (4.21). The last term there (proportional to $\sin \gamma$) involves the commutator function, which vanishes for spacelike separated points.

By analytic continuation then—in fact, precisely the analytic continuation that allowed us to match the $\cos \gamma$ term—we find that

$${}_{\alpha\gamma} \langle \psi | \mathcal{O}_1(t, \Omega) \mathcal{O}_2(t', \Omega') | \psi \rangle_{\alpha\gamma} \tag{7.9}$$

also matches the bulk propagator between the two boundaries. However, direct computation shows agreement only for $\gamma = 0, \pi$. (For example, the coefficient of $\sinh^2 \alpha$ involves $e^{\pm 2i\gamma}$ instead of being γ -independent.) We suspect that this is related to the CPT noninvariance of the $\gamma \neq 0, \pi$ vacua.

C. Entropy

The “CFT calculation” (6.15) now shows that the entropy of the bulk is α -dependent. Thus, black hole entropy need not be $A/4$! It would be interesting to understand this better.

In particular, this result for the black hole entropy assumes first of all that an equilibrium calculation of the entropy holds for these nonequilibrium (but steady state) α -vacua. This need not be true.

Additionally, naïve application of the equilibrium thermodynamic formula

$$\frac{dS}{dE} = \beta, \tag{7.10}$$

leads to the fairly nonsensical, energy-dependent temperature¹²

$$\beta_{\alpha\gamma}(E_i) = -\ln \left| \frac{e^{-\frac{\beta E_i}{2}} \cosh \alpha + e^{i\gamma} \sinh \alpha}{\cosh \alpha + e^{-\frac{\beta E_i}{2} + i\gamma} \sinh \alpha} \right|^2. \quad (7.11)$$

Again, the obvious resolution is that the standard equilibrium formulas do not apply in this nonequilibrium situation. This leaves as an open question whether the black hole entropy is, or is not, $A/4$ for all α , but this is a question that can be addressed using the dual CFT.

Acknowledgments

It is a pleasure to thank Sumit Das, Moshe Elitzur, Gary Gibbons, Finn Larsen, Andreas Karch, Don Marolf, Matt Martin, Robert McNees, Emil Mottola, Simon Ross, Al Shapere, Anastasia Volovich and Xinkai Wu for useful conversations. This work was supported in part by National Science Foundation grant Nos PHY-0244811 and PHY-0555444 and by a Department of Energy contract No. DE-FG01-00ER45832. At The Ohio State University, J.M. is supported in part by Department of Energy contract No. DE-FG02-91ER-40690.

APPENDIX A: DERIVATION OF THE CFT DENSITY MATRIX

In this appendix we derive the density matrix (6.14). Standard arguments [3, 6] and references therein, allow us to write

$$|\tilde{0}\rangle_1 |\tilde{0}\rangle_2 = \text{sech } \alpha e^{e^{-i\gamma} \tanh \alpha a_1^\dagger a_2^\dagger} |0\rangle_1 |0\rangle_2. \quad (A.1)$$

Now we state the main formula we need, which is a version of [69, (A5.17)]. Suppose we have a set of operators $K_\pm, K_3$ obeying

$$[K_3, K_\pm] = \pm K_\pm, \quad [K_+, K_-] = -2K_3. \quad (A.2)$$

We will use

$$K_+ = a_1^\dagger a_2^\dagger, \quad K_- = a_1 a_2, \quad K_3 = \frac{1}{2}(a_1^\dagger a_1 + a_2 a_2^\dagger), \quad (A.3)$$

(note the ordering in K_3). Then, for c-numbers $\gamma_\pm, \gamma_3$ related by

$$\frac{1}{4}\gamma_3^2 = \gamma_+ \gamma_-, \quad (A.4)$$

and setting,

$$\Gamma_\pm = \frac{\gamma_\pm}{1 - \frac{\gamma_3}{2}}, \quad \Gamma_3 = \frac{1}{\left(1 - \frac{\gamma_3}{2}\right)^2}, \quad (A.5)$$

¹² But note that plugging this into the Boltzmann formula does in fact reproduce Eq. (4.44)!

one has the identity,

$$\exp[\gamma_+ K_+ - \gamma_3 K_3 + \gamma_- K_-] = \exp[\Gamma_+ K_+] \exp[(\log \Gamma_3) K_3] \exp[\Gamma_- K_-]. \quad (\text{A.6})$$

We apply this first to

$$\exp\left[e^{-\frac{\beta}{2}} b_1^\dagger b_2^\dagger\right] = \exp\left[e^{-\frac{\beta}{2}} \cosh^2 \alpha K_+ - 2e^{-\frac{\beta}{2}} e^{i\gamma} \cosh \alpha \sinh \alpha K_3 + e^{-\frac{\beta}{2}} e^{2i\gamma} \sinh^2 \alpha K_-\right]. \quad (\text{A.7})$$

so that,

$$|\psi\rangle_{\alpha\gamma} = \text{sech } \alpha \sqrt{1 - e^{-\beta}} e^{\Gamma_+ a_1^\dagger a_2^\dagger} e^{(\log \Gamma_3) K_3} e^{\Gamma_- a_1 a_2} e^{e^{-i\gamma} \tanh \alpha a_1^\dagger a_2^\dagger} |0\rangle_1 |0\rangle_2, \quad (\text{A.8})$$

We can now commute the right-most factor to the left, though this requires another application of (A.6). This gives,

$$|\psi\rangle_{\alpha\gamma} = \frac{\text{sech } \alpha \sqrt{1 - e^{-\beta}}}{1 + e^{-\frac{\beta}{2} + i\gamma} \tanh \alpha} \exp\left[\frac{e^{-\frac{\beta}{2}} + e^{-i\gamma} \tanh \alpha}{1 + e^{-\frac{\beta}{2} + i\gamma} \tanh \alpha} a_1^\dagger a_2^\dagger\right] |0\rangle_1 |0\rangle_2, \quad (\text{A.9})$$

which immediately implies Eq. (6.14).

As a (trivial) check, one can see that setting $\alpha = 0$ reproduces the standard density matrix, and that in general $\text{Tr } \rho_{\alpha\gamma} = 1$.

APPENDIX B: STRINGS ON RINDLER SPACE

In this section, we elaborate on the claims made in §V D that strings do not suffer from pinch singularities in Rindler space. Many of the field theory details we provide below can be found in [46], but are repeated here to set notation and for extension to string theory.

1. The Setup and Vertex Operators

Consider the d -dimensional Rindler spacetime with metric

$$ds^2 = e^{2a\xi} (-d\eta^2 + d\xi^2) + d\vec{y}^2. \quad (\text{B.1})$$

Here a is a constant, η is time, ξ is a spatial coordinate and $\vec{y}$ are $d - 2$ additional, flat, spatial coordinates. There is a horizon at $\xi = -\infty$; $\xi = +\infty$ is spatial infinity. Indeed, the geometry becomes somewhat more illuminating via the coordinate transformation

$$t = a^{-1} e^{a\xi} \sinh(a\eta), \quad x = a^{-1} e^{a\xi} \cosh(a\eta), \quad (\text{B.2})$$

which brings the metric to the form

$$ds^2 = -dt^2 + dx^2 + d\vec{y}^2, \quad (\text{B.3})$$

i.e. Minkowski space, for Rindler space is the spacetime seen by an accelerating observer in a Minkowski background.

Rindler space is plotted in Fig. 9. The region R is that covered by the coordinates (B.1). The region L is the other Rindler region corresponding to $t \rightarrow -t$ and $x \rightarrow -x$ in (B.2),




FIG. 9: Rindler space. R is the Rindler region, L is the opposite Rindler region, F is the future and P is the past. The direction of time (η) in each region is also shown. The diagonal lines are the past and future Rindler horizons. Each point depicts an $\mathbb{R}^{d-2}$.

or equivalently, $\text{Im } \eta = -\pi i a^{-1}$. Other regions not covered by the coordinates (B.1) are the future (F) and past (P). These correspond to $\text{Im } \xi = \frac{\pi}{2a}$ with respectively $\text{Im } \eta = -\frac{\pi}{2a}$, and $\text{Im } \eta = -\frac{3\pi}{2a}$. Equivalently, the other regions are covered via

$$t = a^{-1} e^{a\xi_F} \cosh(a\eta_F), \quad x = a^{-1} e^{a\xi_F} \sinh(a\eta_F), \quad (\text{B.4})$$

$$t = -a^{-1} e^{a\xi_P} \cosh(a\eta_P), \quad x = -a^{-1} e^{a\xi_P} \sinh(a\eta_P), \quad (\text{B.5})$$

$$t = -a^{-1} e^{a\xi_L} \sinh(a\eta_L), \quad x = -a^{-1} e^{a\xi_L} \cosh(a\eta_L), \quad (\text{B.6})$$

which can be summarized via

$$\eta = \eta_L - \frac{\pi}{a}i, \xi = \xi_L, \quad \eta = \eta_F - \frac{\pi}{2a}i, \xi = \xi_F + \frac{\pi}{2a}i, \quad \eta = \eta_P - \frac{3\pi}{2a}i, \xi = \xi_P + \frac{\pi}{2a}i, \quad (\text{B.7})$$

Note the periodicity of (B.2) under $\eta \rightarrow \eta - 2\pi i a^{-1}$. Also, note that “time translation”, defined with respect to η , goes backwards in region L and is spacelike in the future and past. This is analogous to a black hole of temperature a .

Now consider a mass μ scalar field on Rindler space. The wave equation reads

$$[e^{-2a\xi}(-\partial_\eta^2 + \partial_\xi^2) + \vec{p}_y^2 - \mu^2] \phi = 0. \quad (\text{B.8})$$

The positive frequency modes are

$$\phi_{\omega, \vec{k}}^R = \sqrt{\frac{\sinh(\frac{\pi\omega}{a})}{a}} \frac{2}{(2\pi)^{d/2}} e^{-i\omega\eta + i\vec{k}\cdot\vec{y}} K_{i\frac{\omega}{a}} \left(\frac{\sqrt{\mu^2 + \vec{k}^2}}{a} e^{a\xi} \right), \quad (\text{B.9})$$

where $K_\nu(z)$ is a modified Bessel function. (The second solution blows up at infinity, and so is excluded.) These are (Klein-Gordon) δ -function normalized,

$$i \int_{-\infty}^{\infty} d\xi \int d^{d-2}\vec{y} \left[\phi_{\omega, \vec{k}}^{R*} \overleftrightarrow{\partial}_\eta \phi_{\omega', \vec{k}'}^R \right] = \delta(\omega - \omega') \delta^{(d-2)}(\vec{k} - \vec{k}'). \quad (\text{B.10})$$

These Rindler modes can be analytically continued into the future and past (Fig. 9) and are defined to vanish in the other (left) Rindler region. One similarly defines positive frequency modes in the left Rindler region.

As pointed out in §VD, the (tachyon) vertex operators of the (bosonic) string are essentially the on-shell field theory modes. In Minkowski space, the vertex operators are (suppressing the normal ordering symbol) $e^{ikX} = e^{-iEt + i\ell x + i\vec{k}\cdot\vec{y}}$, with $k^2 = -\mu^2 = \frac{1}{\alpha'}$ for the open string. For the right Rindler region, we can use the formula¹³

$$\begin{aligned} \mathcal{V}_{\omega, \vec{k}}^R &= \sqrt{\frac{\sinh\left(\frac{\pi\omega}{a}\right)}{a}} \frac{2}{(2\pi)^{d/2}} e^{-i\omega\eta + i\vec{k}\cdot\vec{y}} K_{i\frac{\omega}{a}}\left(\frac{\sqrt{\mu^2 + \vec{k}^2}}{a} e^{a\xi}\right) \\ &= \frac{1}{2(2\pi)^{d/2} \sqrt{a \sinh \frac{\pi\omega}{a}}} \int_{-\infty}^{\infty} \frac{d\ell}{\sqrt{\mu^2 + \vec{k}^2 + \ell^2}} \left\{ e^{\frac{\pi\omega}{2a} - i\frac{\omega}{a} \sinh^{-1} \frac{\ell}{\sqrt{\mu^2 + \vec{k}^2}} - i\sqrt{\mu^2 + \ell^2 + \vec{k}^2} t + i\ell x + i\vec{k}\cdot\vec{y}} \right. \\ &\quad \left. - e^{-\frac{\pi\omega}{2a} + i\frac{\omega}{a} \sinh^{-1} \frac{\ell}{\sqrt{\mu^2 + \vec{k}^2}} + i\sqrt{\mu^2 + \ell^2 + \vec{k}^2} t + i\ell x + i\vec{k}\cdot\vec{y}} \right\}, \end{aligned} \quad (\text{B.11})$$

to write the on-shell Rindler modes in terms of on-shell Minkowski ones. Thus, the right-hand side of (B.11) is a well-defined expression for the tachyon vertex operator in the right Rindler region.

The left Rindler region vertex operators are similarly

$$\begin{aligned} \mathcal{V}_{\omega, \vec{k}}^L &= \frac{1}{2(2\pi)^{d/2} \sqrt{a \sinh \frac{\pi\omega}{a}}} \int_{-\infty}^{\infty} \frac{d\ell}{\sqrt{\mu^2 + \vec{k}^2 + \ell^2}} \left\{ e^{\frac{\pi\omega}{2a} + i\frac{\omega}{a} \sinh^{-1} \frac{\ell}{\sqrt{\mu^2 + \vec{k}^2}} - i\sqrt{\mu^2 + \ell^2 + \vec{k}^2} t + i\ell x + i\vec{k}\cdot\vec{y}} \right. \\ &\quad \left. - e^{-\frac{\pi\omega}{2a} - i\frac{\omega}{a} \sinh^{-1} \frac{\ell}{\sqrt{\mu^2 + \vec{k}^2}} + i\sqrt{\mu^2 + \ell^2 + \vec{k}^2} t + i\ell x + i\vec{k}\cdot\vec{y}} \right\}. \end{aligned} \quad (\text{B.12})$$

Having written the vertex operators in terms of Minkowski ones, it is now a relatively simple matter to compute correlation functions from the Minkowski space ones.

2. Rindler Space α -Vacua

It is convenient to define the linear combinations of left and right Rindler region modes

$$\begin{aligned} \phi_{\omega, \vec{k}}^1 &= \frac{1}{\sqrt{2}} \left(\phi_{\omega, \vec{k}}^R + i\phi_{\omega, \vec{k}}^L \right), \\ \phi_{\omega, \vec{k}}^2 &= \frac{1}{\sqrt{2}} \left(\phi_{\omega, \vec{k}}^L + i\phi_{\omega, \vec{k}}^R \right). \end{aligned} \quad (\text{B.13})$$

These modes are orthonormal—and, in particular, ϕ^1 are orthogonal to ϕ^2 —if the Rindler modes are. As these combinations only involve positive frequencies, the vacuum defined by the modes (B.13) is again the Rindler vacuum.

¹³ In addition to suppressing the normal ordering symbols, we also suppress the worldsheet dependence of the vertex operator which appears via the string coordinates t , x and $\vec{y}$.




FIG. 10: The 1-loop contribution to the three point function in (a) field theory and (b) the planar contribution to string theory. As position-space diagrams, the pinch singularities appear for α -vacua in field theory, when, say, both the integrated points x and y , and y and z are null-separated, so that the y -integration has a double pole, with potentially conflicting $i\epsilon$ prescriptions. The string theory diagram shows that this is smoothed out.

Now consider the modes

$$\begin{aligned}\phi_{\omega,\vec{k}}^{1(\alpha,\beta)} &= \cosh \alpha \phi_{\omega,\vec{k}}^1 + e^{i\beta} \sinh \alpha \phi_{\omega,-\vec{k}}^{1*}, \\ \phi_{\omega,\vec{k}}^{2(\alpha,\beta)} &= \cosh \alpha \phi_{\omega,\vec{k}}^2 + e^{i\beta} \sinh \alpha \phi_{\omega,-\vec{k}}^{2*},\end{aligned}\tag{B.14}$$

Again, these modes are orthonormal. These modes define the α -vacuum $|\alpha, \beta\rangle$.

It must be emphasized that unlike the black hole α -vacua, the identification (*cf.* §II) associated with this definition of Rindler space α -vacua is afflicted with a fixed point at the origin of Minkowski space.

3. No Pinch Singularities in One-loop Diagrams

Let us now consider one-loop diagrams. For simplicity, we consider bosonic open string diagrams with external tachyon legs, and recall [70] that the mass-squared of the tachyon is $\mu^2 = -\frac{1}{\alpha'}$. The field theory diagram, eq. (5.11), suffered from a pinch singularity. The open string theory analog of this diagram is depicted in Fig. 6 (page 23). It is expected to be nonsingular; in particular, it is equivalent to a tree-diagram in which a single closed string is exchanged. However, the closed string picture is good for the regime of modular parameter which is opposite to the field theory limit. Thus, the mechanism in which the pinch singularity is resolved is inherently stringy.

The simple analysis performed here will be complicated by the fact that string theory diagrams are necessarily on-shell. Thus, for example, the two-point function which epitomized the pinch singularity in field theory is not easy to analyze in string theory—the propagator is by its very nature singular on-shell. So we will examine the three-point function at one-loop, Fig. 10, and see that it also does not appear to have problems that signal a pinch singularity. Rather, we will see how the pinch singularity is regulated.

a. Minkowski space diagrams

We will need Minkowski space one-loop amplitudes. The general result is well-known [70]; we have adapted it from [57]. In terms of the open-string modular parameter, t , the ratio

between the radius and the length of the cylinder, and the vertex operator positions τ , one finds that the M -point amplitude with the first N vertex operators on one boundary and the remaining $M - N$ on the other boundary, is

$$\begin{aligned} \mathcal{A}(k_1, \dots, k_N; k_{N+1}, \dots, k_M) \\ = ig^M \delta^{(26)} \left(\sum_{p=1}^M k_p \right) \int_0^\infty \frac{dt}{2t} (8\pi^2 \alpha' t)^{-13} \eta(it)^{-24} \left[\prod_{p=1}^M \int_0^{2\pi t} d\tau_p \right] \Psi_1 \Psi_2 \Psi_{12}, \end{aligned} \quad (\text{B.15})$$

where

$$\begin{aligned} \Psi_1 &= \prod_{1 \leq i < j \leq N} |\psi_{ij}|^{2\alpha' k_i \cdot k_j}, \quad \Psi_2 = \prod_{N+1 \leq r < s \leq M} |\psi_{rs}|^{2\alpha' k_r \cdot k_s}, \quad \Psi_{12} = \prod_{\substack{1 \leq i \leq N \\ N+1 \leq r \leq M}} (\psi_{ir}^T)^{2\alpha' k_i \cdot k_r}, \\ \psi_{ij} &= -2\pi i \exp\left(-\frac{\tau_{ij}^2}{4\pi t}\right) \frac{\vartheta_{11}\left(i\frac{\tau_{ij}}{2\pi}|it\right)}{\vartheta'_{11}(0|it)}, \quad \psi_{ir}^T = 2\pi \exp\left(-\frac{\tau_{ir}^2}{4\pi t}\right) \frac{\vartheta_{10}\left(i\frac{\tau_{ir}}{2\pi}|it\right)}{\vartheta'_{11}(0|it)}, \end{aligned} \quad (\text{B.16})$$

$\tau_{ij} = \tau_i - \tau_j$, ϑ and η are Jacobi ϑ - and Dedekind η -functions using the conventions of [70], and g is the open string coupling.

b. The Three-Point Function

A quantity which we expect to contribute to a pinch singularity is the one-loop contribution to $\left\langle \phi_{\omega_1, \vec{k}_1}^R \phi_{\omega_2, \vec{k}_2}^R \phi_{\omega_3, \vec{k}_3}^L \right\rangle$. This is given, up to factors of 2, π , and g , by

$$\begin{aligned} \mathcal{A}_{RRL} &= \frac{1}{\sqrt{\sinh \frac{\pi\omega_1}{a} \sinh \frac{\pi\omega_2}{a} \sinh \frac{\pi\omega_3}{a}}} \int_{-\infty}^\infty \frac{d\ell_1}{\sqrt{\mu_1^2 + \ell_1^2}} \int_{-\infty}^\infty \frac{d\ell_2}{\sqrt{\mu_2^2 + \ell_2^2}} \int_{-\infty}^\infty \frac{d\ell_3}{\sqrt{\mu_3^2 + \ell_3^2}} \\ &\times \sum_{\sigma_1, \sigma_2, \sigma_3 = \pm} \sigma_1 \sigma_2 \sigma_3 e^{\frac{\pi}{2a}(\sigma_1 \omega_1 + \sigma_2 \omega_2 + \sigma_3 \omega_3)} e^{-i\sigma_1 \frac{\omega_1}{a} \sinh^{-1} \frac{\ell_1}{\mu_1} - i\sigma_2 \frac{\omega_2}{a} \sinh^{-1} \frac{\ell_2}{\mu_2} + i\sigma_3 \frac{\omega_3}{a} \sinh^{-1} \frac{\ell_3}{\mu_3}} \\ &\times \delta\left(\sigma_1 \sqrt{\mu_1^2 + \ell_1^2} + \sigma_2 \sqrt{\mu_2^2 + \ell_2^2} + \sigma_3 \sqrt{\mu_3^2 + \ell_3^2}\right) \delta(\ell_1 + \ell_2 + \ell_3) \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \\ &\times \int_0^\infty \frac{dt}{2t} (8\pi \alpha' t)^{-13} \eta(it)^{-24} \int_0^{2\pi t} d\tau_1 \int_0^{2\pi t} d\tau_2 \int_0^{2\pi t} d\tau_3 e^{-\frac{\alpha'}{2\pi t} \sum_{i < j} k_i \cdot k_j \tau_{ij}^2} \\ &\times \left| 2\pi \frac{\vartheta_{1\cdot}\left(i\frac{\tau_{12}}{2\pi}|it\right)}{\vartheta'_{11}(0|it)} \right|^{2\alpha' k_1 \cdot k_2} \left| 2\pi \frac{\vartheta_{1\cdot}\left(i\frac{\tau_{13}}{2\pi}|it\right)}{\vartheta'_{11}(0|it)} \right|^{2\alpha' k_1 \cdot k_3} \left| 2\pi \frac{\vartheta_{1\cdot}\left(i\frac{\tau_{23}}{2\pi}|it\right)}{\vartheta'_{11}(0|it)} \right|^{2\alpha' k_2 \cdot k_3}. \end{aligned} \quad (\text{B.17a})$$

where the σ_i are associated with the signs in the exponentials in the two terms of (B.11) and (B.12); the missing index in the ϑ -functions is 1 or 0 depending on whether the associated vertex operators are on the same or different boundaries; it is understood that $k_i = (\sigma_i \sqrt{\mu_i^2 + \ell_i^2}, \ell_i, \vec{k}_i)$; and we have made the convenient definition

$$\mu_i = \sqrt{\mu^2 + \vec{k}_i^2}, \quad \mu^2 = -\frac{1}{\alpha'}, \quad (\text{B.17b})$$

in terms of the spatial momentum $\vec{k}_i$ at each vertex, and the tachyon mass. In fact, the integrals are simplified by using the effective kinematics associated with a 1+1-dimensional 3-particle scattering for particles of mass μ_i and momentum ℓ_i .

The energy-momentum conserving δ -functions imply that one of the σ 's must be different from the other two. Consider the term for which the $\sigma_1 = \sigma_2 = -\sigma_3$; we will obtain the other terms by permutation. We then use the δ -functions to integrate over and eliminate ℓ_1 and ℓ_2 . The ℓ_3 integration is simplified by defining $\zeta = \sinh^{-1} \frac{\ell_3}{\sigma_3 \mu_3}$, which is the boost parameter required to transform the kinematics from the $\ell_3 = 0$ frame of the effective 1+1-dimensional scattering problem. One then finds that the δ -functions impose

$$\ell_1 = \mu_1 \sinh(\sigma_1 \zeta \pm \lambda_1), \quad \ell_2 = \mu_2 \sinh(\sigma_2 \zeta \mp \lambda_2), \quad (\text{B.18a})$$

where the arbitrary sign is the 1 + 1-dimensional analogue of the arbitrary angle which appears in higher-dimensional scattering, and

$$\lambda_1 \equiv \sinh^{-1} \frac{\kappa^2}{2\mu_1\mu_3}, \quad \lambda_2 \equiv \sinh^{-1} \frac{\kappa^2}{2\mu_2\mu_3}, \quad \kappa^4 \equiv \mu_1^4 + \mu_2^4 + \mu_3^4 - 2\mu_1^2\mu_2^2 - 2\mu_1^2\mu_3^2 - 2\mu_2^2\mu_3^2. \quad (\text{B.18b})$$

Therefore,

$$k_1 \cdot k_2 = -\mu_1\mu_2 \cosh(\lambda_1 + \lambda_2) + \vec{k}_1 \cdot \vec{k}_2, \quad (\text{B.19a})$$

$$k_1 \cdot k_3 = \mu_1\mu_3 \cosh \lambda_1 + \vec{k}_1 \cdot \vec{k}_3, \quad (\text{B.19b})$$

$$k_2 \cdot k_3 = \mu_2\mu_3 \cosh \lambda_2 + \vec{k}_2 \cdot \vec{k}_3, \quad (\text{B.19c})$$

which is independent of both ζ and the sign in (B.18a). The ℓ_3 —or rather ζ —integration then yields the expected ω -constraining δ -function; the final result, upon summing over the arbitrary sign in (B.18a), is thus

$$\begin{aligned} \mathcal{A}_{RRL} = & \frac{1}{\sqrt{\sinh \frac{\pi\omega_1}{a} \sinh \frac{\pi\omega_2}{a} \sinh \frac{\pi\omega_3}{a}}} \delta(\omega_1 + \omega_2 - \omega_3) \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \\ & \times \left\{ 2 \frac{\sinh \left[\frac{\pi}{2a} (\omega_1 + \omega_2 - \omega_3) \right] \cos(\lambda_1 - \lambda_2)}{\mu_1\mu_2 \sinh(\lambda_1 + \lambda_2)} \int_0^\infty \frac{dt}{2t} (8\pi\alpha't)^{-13} \eta(it)^{-24} \right. \\ & \times \int_0^{2\pi t} d\tau_1 \int_0^{2\pi t} d\tau_2 \int_0^{2\pi t} d\tau_3 e^{-\frac{\alpha'}{2\pi t} k_1 \cdot k_2 \tau_{12}^2} \left| 2\pi \frac{\vartheta_{1, \left(i \frac{\tau_{12}}{2\pi} | it \right)}}{\vartheta'_{11}(0|it)} \right|^{2\alpha' k_1 \cdot k_2} \\ & \times e^{-\frac{\alpha'}{2\pi t} k_1 \cdot k_3 \tau_{13}^2} \left| 2\pi \frac{\vartheta_{1, \left(i \frac{\tau_{13}}{2\pi} | it \right)}}{\vartheta'_{11}(0|it)} \right|^{2\alpha' k_1 \cdot k_3} e^{-\frac{\alpha'}{2\pi t} k_2 \cdot k_3 \tau_{23}^2} \left| 2\pi \frac{\vartheta_{1, \left(i \frac{\tau_{23}}{2\pi} | it \right)}}{\vartheta'_{11}(0|it)} \right|^{2\alpha' k_2 \cdot k_3} \\ & \left. + 2 \text{ perms} \right\}, \quad (\text{B.20}) \end{aligned}$$

where $k_i \cdot k_j$ are given in terms of the $\vec{k}$'s and the masses by (B.19). The sign in the energy conserving δ -function is due to the leftness of the Rindler mode, not the relative value of σ_3 , but the sign in the $\sinh$ in the numerator is due to the relative value of σ_3 . Because the λ 's and μ 's depend only on the $\vec{k}$'s, and appear in the amplitude (B.20) via $k_i \cdot k_j$ for on-shell k 's, the final integrations to be performed are identical to those for Minkowski space amplitudes, which are well-known to be well-behaved. Thus we see that the stringy amplitudes for α -vacua in Rindler space are also well-behaved.

i. Open String Factorization It is the large t regime of (B.20) from which the field theory pinch singularities are expected. Approximating the integrand of the planar diagram at large t , and, using the translational symmetry of the annulus to fix τ_1 to a convenient value yields, (cf. e.g. [57])

$$\begin{aligned} \mathcal{A}_{RRL} = & \frac{1}{\sqrt{\sinh \frac{\pi\omega_1}{a} \sinh \frac{\pi\omega_2}{a} \sinh \frac{\pi\omega_3}{a}}} \delta(\omega_1 + \omega_2 - \omega_3) \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) (1 - y_1)^2 \\ & \times \left\{ 2\pi \frac{\sinh \left[\frac{\pi}{2a} (\omega_1 + \omega_2 - \omega_3) \right] \cos(\lambda_1 - \lambda_2)}{\mu_1 \mu_2 \sinh(\lambda_1 + \lambda_2)} \int_0^\infty dt (8\pi\alpha' t)^{-13} e^{-2\pi t} \right. \\ & \times \left. \int_{-y_0}^{y_0} dy_2 \int_{-y_0}^{y_0} dy_3 e^{-\frac{\alpha'}{2\pi t} \sum_{i < j} k_i \cdot k_j \tau_{ij}^2} |y_{12}|^{2\alpha' k_1 \cdot k_2} |y_{13}|^{2\alpha' k_1 \cdot k_3} |y_{23}|^{2\alpha' k_2 \cdot k_3} + 2 \text{ perms} \right\}, \quad (\text{B.21a}) \end{aligned}$$

where

$$\tau_i = -\ln \left| \frac{1 + y_i}{1 - y_i} \right| + \pi t, \quad y_0 = \tanh \frac{\pi t}{2}, \quad (\text{B.21b})$$

The difference between the planar and the nonplanar diagram, in this approximation, is that the y 's on the “other” boundary would lie in $(-\infty, -y_0^{-1}) \cup (y_0^{-1}, \infty)$ instead of $(-y_0, y_0)$. In fact, in the $t \rightarrow \infty$ approximation, $y_0 \rightarrow 1$. Incorporating this, and introducing a new integration for reasons to become clear, the amplitude (B.21a) can be rewritten as

$$\begin{aligned} \mathcal{A}_{RRL} = & \frac{1}{\sqrt{\sinh \frac{\pi\omega_1}{a} \sinh \frac{\pi\omega_2}{a} \sinh \frac{\pi\omega_3}{a}}} \delta(\omega_1 + \omega_2 - \omega_3) \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) (1 - y_1^2) \\ & \times \left\{ 2 \frac{\sinh \left[\frac{\pi}{2a} (\omega_1 + \omega_2 - \omega_3) \right] \cos(\lambda_1 - \lambda_2)}{\mu_1 \mu_2 \sinh(\lambda_1 + \lambda_2)} \int_0^1 \frac{dq}{2q} \int \frac{d^{26}k}{(2\pi)^{26}} q^{\alpha'(k^2 + \mu^2)} \int_{-1}^1 dy_2 \int_{-1}^1 dy_3 \right. \\ & \times \left. e^{-2\alpha' k \cdot (k_1 \tau_1 + k_2 \tau_2 + k_3 \tau_3)} |y_{12}|^{2\alpha' k_1 \cdot k_2} |y_{13}|^{2\alpha' k_1 \cdot k_3} |y_{23}|^{2\alpha' k_2 \cdot k_3} + 2 \text{ perms} \right\}, \quad (\text{B.22}) \end{aligned}$$

where $q = e^{-2\pi t}$. Integration of the Gaussian integral k yields the original expression (B.21a). However, the form (B.22), shows the open string factorization illustrated in Fig. 7 (page 24). In particular, y_i 's are the vertex operator positions on the upper half-plane and $\pm k$ are the momenta of the extra vertex operator insertions. (The fixing of $y_1 = 0$, as well as the fixing of the extra vertex operator positions at $y = \pm 1$ is maximal fixings allowed by the conformal Killing vectors of the upper half-plane.)

If one integrates q instead of k in (B.22), a factor of $\frac{1}{k^2 + \mu^2}$, the field theory propagator of the intermediate state, is obtained. Moreover, there is a pole that involves $k + k_3$ via the factors of $e^{-2\alpha' k \cdot k_3 (\tau_3 - \pi t)} = \left| \frac{1 + y_3}{1 - y_3} \right|^{-\alpha' (k + k_3)^2 + \alpha' k^2 - 1}$ (where all τ_i 's could be replaced by $\tau_i - \pi t$ due to energy-momentum conservation, and using $\alpha' \mu^2 = -1$). Thus, exchange of a tachyon ($k^2 = -\mu^2 = \frac{1}{\alpha'}$) leads to simultaneous poles which, because the second pole is associated with y_3 —i.e. the term from the left Rindler region which therefore has the opposite $i\epsilon$ prescription—gives the pinch singularity. However, we have already seen from the complete result (B.20) that these apparent pinch singularities smoothed out by the full string theory. This is because the y_3 integration only goes to $\pm y_0$, not ± 1 where the singularity appears.

[1] E. Mottola, *Particle creation in de Sitter space*, Phys. Rev. D **31** (1985) 754–766.

- [2] B. Allen, *Vacuum states in de Sitter space*, Phys. Rev. D **32** (1985) 3136–3149.
- [3] R. Bousso, A. Maloney and A. Strominger, *Conformal vacua and entropy in de Sitter space*, Phys. Rev. D **65** (2002) 104039; [[hep-th/0112218](#)].
- [4] N. Kaloper, M. Kleban, A. Lawrence, S. Shenker and L. Susskind, *Initial Conditions for Inflation*, J. High Energy Phys. **11** (2002) 037; [[hep-th/0209231](#)].
- [5] M. B. Einhorn and F. Larsen, *Interacting quantum field theory in de Sitter vacua*, Phys. Rev. D **67** (024001) 2003; [[hep-th/0209159](#)].
- [6] M. B. Einhorn and F. Larsen, *Squeezed states in the de Sitter vacuum*, Phys. Rev. D **68** (2003) 064002; [[hep-th/0305056](#)].
- [7] K. Goldstein and D. A. Lowe, *A note on alpha-vacua and interacting field theory in de Sitter space*, Nucl. Phys. **B669** (2003) 325; [[hep-th/0302050](#)].
- [8] K. Goldstein and D. A. Lowe, *Real-time perturbation theory in de Sitter space*, Phys. Rev. D **69** (2004) 023507; [[hep-th/0308135](#)].
- [9] R. Brunetti, K. Fredenhagen and S. Hollands, *A remark on alpha vacua for quantum field theories on de Sitter space*, J. High Energy Phys. **05** (2005) 063; [[hep-th/0503022](#)].
- [10] H. Collins and R. Holman, *Taming the alpha vacuum*, Phys. Rev. D **70** (2004) 084019; [[hep-th/0312143](#)].
- [11] S. Naidu and R. Holman, *Inflaton decay in an alpha vacuum*, Phys. Rev. D **71** (2005) 064026; [[hep-th/0409013](#)].
- [12] H. Collins, R. Holman and M. R. Martin *The Fate of the Alpha-Vacuum*, Phys. Rev. D **68** (2003) 124012; [[hep-th/0306028](#)].
- [13] H. Collins and M. R. Martin, *The enhancement of inflaton loops in an alpha-vacuum*, Phys. Rev. D **70** (2004) 084021; [[hep-ph/0309265](#)].
- [14] R. Easther, B. R. Greene, W. H. Kinney and G. Shiu *Inflation as a Probe of Short Distance Physics*, Phys. Rev. D **64** (2001) 103502; [[hep-th/0104102](#)].
- [15] R. Easther, B. R. Greene, W. H. Kinney and G. Shiu *Imprints of Short Distance Physics On Inflationary Cosmology*, Phys. Rev. D **67** (2003) 063508; [[hep-th/0110266](#)].
- [16] U. H. Danielsson, *A note on inflation and transplanckian physics*, Phys. Rev. D **66** (2002) 023511; [[hep-th/0203198](#)].
- [17] U. H. Danielsson, *Inflation, holography, and the choice of vacuum in de Sitter space*, J. High Energy Phys. **07** (2002) 040; [[hep-th/0205227](#)].
- [18] J. de Boer, V. Jejjala and D. Minic, *Alpha-states in de Sitter Space*, Phys. Rev. D **71** (2005) 044013; [[hep-th/0406217](#)].
- [19] B. R. Greene, K. Schalm, G. Shiu and J. P. van der Schaar, *Decoupling in an expanding universe: backreaction barely constrains short distance effects in the CMB*, J. Cosmol. Astropart. Phys. **02** (2005) 001; [[hep-th/0411217](#)].
- [20] J. Martin and R. H. Brandenberger, *The Trans-Planckian Problem of Inflationary Cosmology*, Phys. Rev. D **63** (2001) 123501; [[hep-th/0005209](#)].
- [21] R. H. Brandenberger and J. Martin, *The Robustness of Inflation to Changes in Super-Planck-Scale Physics*, Mod. Phys. Lett. **A6** (2001) 999–1006; [[astro-ph/0005432](#)].
- [22] R. H. Brandenberger, S. E. Joras and J. Martin, *Trans-Planckian Physics and the Spectrum of Fluctuations in a Bouncing Universe*, Phys. Rev. D **66** (2002) 083514; [[hep-th/0112122](#)].
- [23] R. H. Brandenberger and J. Martin, *On Signatures of Short Distance Physics in the Cosmic Microwave Background*, Int. J. Mod. Phys. **A17** (2002) 3663–3680; [[hep-th/0202142](#)].
- [24] F. Larsen and R. McNees, *Inflation and de Sitter Holography*, J. High Energy Phys. **07** (2003) 051; [[hep-th/0307026](#)].

- [25] H. Collins, *Fermionic alpha-vacua*, Phys. Rev. D **71** (2005) 024002; [hep-th/0410229].
- [26] Juan M. Maldacena, *Eternal Black Holes in AdS*, J. High Energy Phys. **04** (2003) 021; [hep-th/0106112].
- [27] V. Balasubramanian, T. S. Levi and A. Naqvi, *A comment on multiple vacua, particle production and the time dependent AdS/CFT correspondence*, Phys. Rev. D **69** (2004) 124003; [hep-th/0303157].
- [28] S. F. Ross and G. Titchener, *Time-dependent spacetimes in AdS/CFT: Bubble and black hole*, J. High Energy Phys. **02** (2005) 021; [hep-th/0411128].
- [29] M. D. Kruskal, *Maximal Extension of Schwarzschild Metric*, Phys. Rev. **119** (1960) 1743.
- [30] G. Gibbons, *The elliptic interpretation of black holes and quantum mechanics*, Nucl. Phys. **B271** (1986) 497–508.
- [31] A. Chamblin and G. Gibbons, *Nucleating black holes via non-orientable instantons*, Phys. Rev. D **55** (1997) 2177–2185; [gr-qc/9607079].
- [32] L. Fidkowski, V. Hubeny, M. Kleban and S. Shenker, *The Black Hole Singularity in AdS/CFT*, J. High Energy Phys. **02** (2004) 014; [hep-th/0306170].
- [33] C. P. Herzog and D. T. Son, *Schwinger-Keldysh propagators from AdS/CFT correspondence*, J. High Energy Phys. **03** (2003) 046; [hep-th/0212072].
- [34] J. B. Hartle and S. W. Hawking, *Path-integral derivation of black-hole radiance*, Phys. Rev. D **13** (1976) 2188.
- [35] M. Bañados, C. Teitelboim and J. Zanelli, *The Black Hole in Three-Dimensional Space-Time*, Phys. Rev. Lett. **69** 19921849–1851; [hep-th/9204099].
- [36] M. Bañados, M. Henneaux, C. Teitelboim and J. Zanelli, *Geometry of the (2+1) Black Hole*, Phys. Rev. D **48** (1993) 1506–1525; [gr-qc/9302012].
- [37] K. Shiraishi and T. Maki, *Quantum Fluctuation of Stress Tensor and Black Holes in Three Dimensions*, Phys. Rev. D **49** (1994) 5286–5294.
- [38] I. Ichinose and Y. Satoh, *Entropies Of Scalar Fields On Three-Dimensional Black Holes*, Nucl. Phys. **B447** (1995) 340; [hep-th/9412144].
- [39] C. Martínez and J. Zanelli, *Back-reaction of a conformal field on a three-dimensional black hole*, Phys. Rev. D **55** (1997) 3642–3646; [gr-qc/9610050].
- [40] E. Keski-Vakkuri, *Bulk and boundary dynamics in BTZ black holes*, Phys. Rev. D **59** (1999) 104001; [hep-th/9808037].
- [41] S. Hemming, E. Keski-Vakkuri and P. Kraus, *Strings in the extended BTZ spacetime*, J. High Energy Phys. **10** (2002) 006; [hep-th/0208003].
- [42] P. Kraus, H. Ooguri and S. Shenker, *Inside the Horizon with AdS/CFT*, Phys. Rev. D **67** (2003) 124022; [hep-th/0212277].
- [43] C. P. Burgess and C. A. Lutken, *Propagators and Effective Potentials in Anti-de Sitter Space*, Phys. Lett. B **153** (1985) 137–141.
- [44] T. Inami and H. Ooguri, *One-Loop Effective Potential in Anti-de Sitter Space*, Prog. Theor. Phys. **73** (1985) 1051–1054.
- [45] E. Witten, *Anti De Sitter Space And Holography*, Adv. Theor. Math. Phys. **2** (1998) 253–291; [hep-th/9802150].
- [46] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, (Cambridge University Press, Cambridge, 1984).
- [47] L. Susskind, L. Thorlacius and J. Uglum, *The Stretched Horizon and Black Hole Complementarity*, Phys. Rev. D **48** (1993) 3743–3761; [hep-th/9306069].
- [48] D. A. Lowe, J. Polchinski, L. Susskind, L. Thorlacius and J. Uglum, *Black Hole Complementarity*, Phys. Rev. D **48** (1993) 3743–3761; [hep-th/9306069].

- tarity* vs. *Locality*, Phys. Rev. D **D52** (1995) 6997–7010; [hep-th/9506138].
- [49] L. Susskind, *Twenty Years of Debate With Stephen*, preprint SU-ITP 02-11 hep-th/0204027.
 - [50] F. Leblond, D. Marolf and R. C. Myers, *Tall tales from de Sitter space I: Renormalization group flows*, J. High Energy Phys. **06** (2002) 052; [hep-th/0202094].
 - [51] R. P. Feynman, *Elementary particles and the laws of physics : the 1986 Dirac memorial lectures*, (Cambridge University Press, Cambridge, 1987).
 - [52] D. Ruelle, *Smooth Dynamics and New Theoretical Ideas in Nonequilibrium Statistical Mechanics*, J. Stat. Phys. **95** (1999) 393–468; [chao-dyn/9812032].
 - [53] M. Elitzur, *Astronomical masers*, (Kluwer Academic Publishers, The Netherlands: 1992), particularly §2.5.
 - [54] H. J. de Vega and N. G. Sanchez, *String Quantization In Accelerated Frames And Black Holes*, Nucl. Phys. **B299** (1988) 818.
 - [55] D. A. Lowe and A. Strominger, *Strings near a Rindler or black hole horizon*, Phys. Rev. D **51** (1995) 1793–1799; [hep-th/9410215].
 - [56] H. Liu, G. Moore and N. Seiberg, *Strings in a Time-Dependent Orbifold*, J. High Energy Phys. **06** (2002) 045; hep-th/0204168.
 - [57] H. Liu and J. Michelson, *Stretched Strings in Noncommutative Field Theory*, Phys. Rev. D **62** (2000) 066003; [hep-th/0004013].
 - [58] J. Polchinski, *Factorization of Bosonic String Amplitudes*, Nucl. Phys. **B307** (1988) 61–92.
 - [59] P. Bedaque, *Thermalization and pinch singularities in non-equilibrium quantum field theory*, Phys. Lett. B **344** (1995) 23–28; [hep-ph/9410415].
 - [60] I. Dadić, *Cancellation of pinching singularities in out-of-equilibrium thermal field theory*, hep-ph/9809395.
 - [61] I. Dadić, *Pinching phenomenon: Central features in out of equilibrium thermal field theories*, hep-ph/0103025.
 - [62] P. Candelas, *Vacuum polarization in Schwarzschild spacetime*, Phys. Rev. D **21** (1980) 2185–2202.
 - [63] I. R. Klebanov and E. Witten, *AdS/CFT Correspondence and Symmetry Breaking*, Nucl. Phys. **B556** (1999) 89–114; [hep-th/9905104].
 - [64] N. P. Landsman and Ch. G. van Weert, *Real- and Imaginary-Time Field Theory at Finite Temperature and Density*, Phys. Rept. **145** (1987) 141–249.
 - [65] V. Balasubramanian, P. Kraus and A. Lawrence, *Bulk vs. Boundary Dynamics in Anti-de Sitter Space-Time*, Phys. Rev. D **59** (1999) 046003; [hep-th/9805171].
 - [66] V. Balasubramanian, P. Kraus, A. Lawrence and S. Trivedi, *Holographic Probes of Anti-de Sitter Space-Times*, Phys. Rev. D **59** (1999) 104021; [hep-th/9808017].
 - [67] T. Banks, M. R. Douglas, G. T. Horowitz and E. Martinec, *AdS Dynamics from Conformal Field Theory*, hep-th/9808016.
 - [68] V. Balasubramanian and S. F. Ross, *Holographic Particle Detection*, Phys. Rev. D **61** (2000) 044007; [hep-th/9906226].
 - [69] S. M. Barnett and P. M. Radmore, *Methods in Theoretical Quantum Optics*, (Oxford University Press, Oxford, 1997).
 - [70] J. Polchinski, *String Theory, Vol. I, An Introduction to the Bosonic String*, (Cambridge University Press, Cambridge, 1998).